


Physicists provide new insights into coral skeleton formation

October 29 2013



Porites Lobata corallite surface: Scanning electron microscopy image of the surface of a Porites sp. coral from which the living tissue (biofilm) has been removed by submersion in fresh water and gentle brushing. The round features are where the polyp mouths were situated. Credit: Renée van de Locht/Dr Roland Kröger

An international team of scientists, led by physicists from the University of York, has shed important new light on coral skeleton formation.


Their investigations, carried out at the nanoscale, provide valuable new information for scientists and environmentalists working to protect and conserve coral from the threats of acidification and rising water temperatures.

As corals grow, they produce limestone – calcium carbonate - skeletons which build up over time into vast reefs. The skeleton's role is to help the living biofilm to move towards the light and nutrients.

Understanding the calcification mechanism by which these skeletons are formed is becoming increasingly important due to the potential impact of climate change on this process.

The scientists looked at the smallest building blocks that can be identified – a microstructure called spherulites – by making a thin cross-section less than 100 nanometres in thickness of a skeleton crystal. They then used Transmission Electron Microscopy (TEM) to analyse the crystals in minute detail.

The TEM micrographs revealed three distinct regions: randomly orientated granular, porous nanocrystals; partly oriented nanocrystals which were more granular and porous; and densely packed aligned needle-like crystals.



Acicular crystals: A) Scanning electron microscopy image of a *Porites lobata* thin section showing the needle-like crystals, inset is a transmission electron microscopy image. B) High resolution transmission electron microscopy of a crystal needle on the nanoscale, showing aragonite crystal lattice. Inset is a fast Fourier transform (FFT) analysis determining the crystal growth direction of the crystal needle. Credit: Renée van de Locht/Dr Roland Kröger

These different regions could be directly correlated to times of the day - at sunset, granular and porous crystals are formed, but as night falls, the calcification process slows down and there is a switch to long aligned needles.

Corresponding author Renée van de Locht, a final-year PhD student with the Department of Physics at the University of York, says, "Coral plays a vital role in a variety of eco-systems and supports around 25 per cent of all marine species. In addition, it protects coastlines from wave erosion and plays a key role in the fisheries and tourism industries. However, the fundamental principles of coral's skeleton formation are still not fully understood.

"It has been suspected for some time that the contrast bands seen in

crystals in optical images were daily bands. Through our research we have been able to show what the crystals actually contain and the differences between day and night crystals."

The research was funded by the Engineering and Physical Sciences Research Council (EPSRC) and the University of York, and concentrated on three species of tropical, reef-building coral - *Porites lobata*, *Siderastrea sidereal*, *Montastrea annularis*.

The experimental work was carried out at the University of York's Department of Physics and the York JEOL Nanocentre, as well as the Centre for Microscopy, Characterisation and Analysis (MCA) at the University of Western Australia.

Lead investigator Dr Roland Kröger says, "Although we knew there was a difference between day and night crystals, we've actually been able to see the evolution from granular to aligned needles and to find out much more information about the phase, orientation and size of the aragonite [crystals](#)."

The York researchers are now turning their attention to looking directly at the affects of acidification. Their latest research studies five-day old coral larvae and compares a population from a normal sea water environment with another in an acidic environment.

The aim is to investigate the nanoscale impacts of the different environments at early growth stage to assess how these could affect the whole colony and the bigger reef.

More information: The first phase of the research was reported in the *Journal of Structural Biology* earlier this year. The article 'Microstructural evolution and nanoscale crystallography in scleractinian coral spherulites' by Renée van de Locht, Andreas Verch, Martin

Saunders, Delphine Dissard, Tim Rixen, Aurélie Moya and Roland Kröger can be seen at: [www.sciencedirect.com/science/ ...
ii/S1047847713001196](http://www.sciencedirect.com/science/.../S1047847713001196)

Provided by University of York

Citation: Physicists provide new insights into coral skeleton formation (2013, October 29)
retrieved 26 April 2024 from
<https://phys.org/news/2013-10-physicists-insights-coral-skeleton-formation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.