


## Simulations help researchers decide which technology would make a better solar collector, quantum dot or nanowire

October 1 2013



Amorphous Silicon nanowire (yellow network) facilitates harvesting of solar energy in the form of a photon (wavy line). In the process of light absorption a pair of mobile charge carriers is created (red clouds depict an electron smeared in space, while the blue clouds visualize the so-called hole which is a positively charged carrier). The energy of their directed motion is then transformed into electricity. Electron and hole charge distributions are often located in different regions of space due to multiple structural defects in amorphous silicon nanowires. Credit: A.Kryjevski, S.Kilina and D.Kilin/JRSE



A trio of researchers at North Dakota State University and the University of South Dakota have turned to computer modeling to help decide which of two competing materials should get its day in the sun as the nanoscale energy-harvesting technology of future solar panels—quantum dots or nanowires.

Andrei Kryjevski and his colleagues, Dimitri Kilin and Svetlana Kilina, report in AIP Publishing's *Journal of Renewable and Sustainable Energy* that they used computational chemistry models to predict the electronic and optical properties of three types of nanoscale (billionth of a meter) silicon structures with a potential application for solar energy collection: a quantum dot, one-dimensional chains of quantum dots and a nanowire. The ability to absorb light is substantially enhanced in nanomaterials compared to those used in conventional semiconductors. Determining which form—quantum dots or nanowire—maximizes this advantage was the goal of the numerical experiment conducted by the three researchers.

"We used Density Functional Theory, a computational approach that allows us to predict electronic and <u>optical properties</u> that reflect how well the nanoparticles can absorb light, and how that effectiveness is affected by the interaction between quantum dots and the disorder in their structures," Kryjevski said. "This way, we can predict how quantum dots, quantum dot chains and nanowires will behave in real life even before they are synthesized and their working properties experimentally checked."

The simulations made by Kryjevski, Kilin and Kilina indicated that light absorption by silicon quantum dot chains significantly increases with increased interactions between the individual nanospheres in the chain. They also found that light absorption by quantum dot chains and <u>nanowires</u> depends strongly on how the structure is aligned in relation to the direction of the photons striking it. Finally, the researchers learned that the atomic structure disorder in the amorphous nanoparticles results



in better <u>light absorption</u> at lower energies compared to crystalline-based nanomaterials.

"Based on our findings, we believe that putting the amorphous quantum dots in an array or merging them into a nanowire are the best assemblies for maximizing the efficiency of silicon nanomaterials to absorb light and transport charge throughout a photovoltaic system," Kryjevski said. "However, our study is only a first step in a comprehensive computational investigation of the properties of semiconductor quantum dot assemblies.

"The next steps are to build more realistic models, such as larger <u>quantum dots</u> with their surfaces covered by organic ligands and simulate the processes that occur in actual solar cells," he added.

**More information:** The article, "Amorphous Silicon Nanomaterials: Quantum Dots Versus Nanowires" by Andrei Kryjevski, Dmitri Kilin and Svetlana Kilina, appears in the *Journal of Renewable and Sustainable Energy*. <u>dx.doi.org/10.1063/1.4817728</u>

## Provided by American Institute of Physics

Citation: Simulations help researchers decide which technology would make a better solar collector, quantum dot or nanowire (2013, October 1) retrieved 21 May 2024 from <u>https://phys.org/news/2013-10-simulations-technology-solar-collector-quantum.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.