


# SMOS sees triple superstorms

October 23 2013



SMOS's microwave radiometer captured wind speed readings from three different typhoons during 10–15 October 2013. The image shows wind speeds up to 140 km/h (dark red) for Cyclone Phailin (left), Typhoon Nari (middle) and Typhoon Wipha (right). Credit: ESA/IFREMER/CLS/CATDS/CNES

ESA's versatile water mission tracked Asia's recent storms over land and sea.

Over the past three weeks, tropical cyclone activity has intensified over the seas bordering southern and eastern Asia.


Cyclone Phailin began forming on 4 October in the Gulf of Thailand,

before moving northwest across the Bay of Bengal and making landfall in northern India. The storm affected about 12 million people and claimed dozens of lives.

Typhoons Nari and Wipha subsequently formed on 8 October: Nari west of the Philippines and Wipha east of Guam. Nari displaced tens of thousands of people in the Philippines and Vietnam, while Wipha lashed Japan's east coast.

During all three superstorms, ESA's Soil Moisture and Ocean Salinity satellite, SMOS, captured snapshots of surface [wind speeds](#) under the intense storms.

SMOS carries a novel microwave sensor to capture images of 'brightness temperature'. These images correspond to radiation emitted from Earth's surface, which are then used to derive information on [soil moisture](#) and ocean salinity.



SMOS reading of soil moisture in southern Asia when Cyclone Phailin hit land in October 2013. Wetter soils are shown in blue with drier soils in yellow and orange, due to the passage of the storm and the intense rain falls it causes. Satellite's ascending and descending orbits are added as time passes, outlined by the red dotted lines.

Strong winds over oceans whip up waves and whitecaps, which in turn affect the microwave radiation from the surface. This means that although strong storms make it difficult to measure salinity, the changes in radiation can, however, be linked directly to the strength of the wind over the sea.

This method of measuring wind speeds, thanks to the fact that SMOS is largely unaffected by atmospheric conditions such as thick clouds or rain, was developed by scientists at France's Ifremer and CLS research centres. It was first used when Hurricane Igor developed over the Atlantic Ocean in 2010.

Being able to measure ocean surface wind in stormy conditions with the broad and frequent coverage that SMOS can provide is paramount for tracking and forecasting storm strength.

"The information that SMOS can provide for storm forecasting has great potential for future operational use of such data. Now that the method has been more consolidated, we are planning to deliver data to the community on a regular basis," says Nicolas Reul from Ifremer.

In addition to tracking this month's storms at sea, SMOS also gathered readings of soil moisture when the storms hit land. The data show which areas experienced the most extreme flooding caused by heaving rainfall

and overflowing rivers. This information could be valuable for early warnings and hydrological forecasting.

Launched in 2009, the versatile SMOS satellite has gone above and beyond its original objectives of mapping soil moisture and [ocean salinity](#). In addition to tracking major storms, its measurements are also providing new information on sea-ice, permafrost and wetlands.

Provided by European Space Agency

Citation: SMOS sees triple superstorms (2013, October 23) retrieved 23 June 2024 from <https://phys.org/news/2013-10-smos-triple-superstorms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.