

# Increasing toxicity of algal blooms tied to nutrient enrichment and climate change

October 24 2013

---



Like pea soup, a thick mat of toxic microcystins cyanobacteria on Lake Taihu in China gets stirred up in the wake of a boat. Credit: Hans Paerl, University of North Carolina

Nutrient enrichment and climate change are posing yet another concern of growing importance: an apparent increase in the toxicity of some algal blooms in freshwater lakes and estuaries around the world, which

threatens aquatic organisms, ecosystem health and human drinking water safety.

As this [nutrient enrichment](#), or "eutrophication" increases, so will the proportion of toxin-producing strains of [cyanobacteria](#) in [harmful algal blooms](#), scientists said.

Researchers from Oregon State University and the University of North Carolina at Chapel Hill will outline recent findings in an analysis Friday in the journal *Science*.

Cyanobacteria are some of the oldest microorganisms on Earth, dating back about 3.5 billion years to a time when the planet was void of oxygen and barren of most life. These bacteria are believed to have produced the oxygen that paved the way for terrestrial life to evolve. They are highly adaptive and persistent, researchers say, and today are once again adapting to new conditions in a way that threatens some of the life they originally made possible.

A particular concern is *Microcystis* sp., a near-ubiquitous cyanobacterium that thrives in warm, nutrient-rich and stagnant waters around the world. Like many cyanobacteria, it can regulate its position in the water column, and often forms green, paint-like scums near the surface.

In a high-light, oxidizing environment, microcystin-producing cyanobacteria have a survival advantage over other forms of cyanobacteria that are not toxic. Over time, they can displace the nontoxic strains, resulting in blooms that are increasingly toxic.



Toxic microcystin bacteria float, along with a dead fish, on the surface of this lake. Credit: Oregon State University

"Cyanobacteria are basically the cockroaches of the aquatic world, they're the uninvited guest that just won't leave," said Timothy Otten, a postdoctoral scholar in the OSU College of Science and College of Agricultural Sciences. His work has been supported by the National Science Foundation.

"When one considers their evolutionary history and the fact that they've persisted even through ice ages and asteroid strikes, it's not surprising they're extremely difficult to remove once they've taken hold in a lake," he said. "For the most part, the best we can do is to try to minimize the conditions that favor their proliferation."

Researchers lack an extensive historical record of bloom events and their associated toxicities to put current observations into a long-term context. However, Otten said, "If you go looking for toxin-producing cyanobacteria, chances are you won't have to look very long until you find some."

There are more than 123,000 lakes greater than 10 acres in size spread across the United States, and based on the last EPA National Lakes Assessment, at least one-third may contain toxin-producing cyanobacteria. Dams; rising temperatures and [carbon dioxide](#) concentrations; droughts; and increased runoff of nutrients from urban and agricultural lands are all compounding the problem.

Many large, eutrophic lakes such as Lake Erie are plagued each year by [algal blooms](#) so massive that they are visible from outer space.

Researchers studying cyanobacterial toxins say it's improbable that their true function was to be toxic, since they actually predate any predators. New research suggests that the potent liver toxin and possible carcinogen, microcystin, has a protective role in cyanobacteria and helps them respond to oxidative stress. This is probably one of the reasons the genes involved in its biosynthesis are so widespread across cyanobacteria and have been retained over millions of years.



Researchers take samples from Lake Taihu in China, when it was heavily contaminated with toxic algal blooms that turned the surface water green. Credit: Oregon State University

Because of their buoyancy and the location of toxins primarily within the cell, exposure risks are greatest near the water's surface, which raises concerns for swimming, boating and other recreational uses.

Also, since cyanobacteria blooms become entrenched and usually occur every summer in impacted systems, chronic exposure to [drinking water](#) containing these compounds is an important concern that needs more attention, Otten said.

"Water quality managers have a toolbox of options to mitigate cyanobacteria toxicity issues, assuming they are aware of the problem and compelled to act," Otten said. "But there are no formal regulations in place on how to respond to bloom events.

"We need to increase public awareness of these issues," he said. "With a

warming climate, rising carbon dioxide levels, dams on more rivers than not, and overloading of nutrients into our waterways, the magnitude and duration of toxic cyanobacterial blooms is only going to get worse."

**More information:** "Blooms Bite the Hand that Feeds Them," by H.W. Paerl et al. *Science*, 2013.

Provided by Oregon State University

Citation: Increasing toxicity of algal blooms tied to nutrient enrichment and climate change (2013, October 24) retrieved 27 April 2024 from <https://phys.org/news/2013-10-toxicity-algal-blooms-tied-nutrient.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--