

# World ocean systems undermined by climate change by 2100

October 15 2013

---

An ambitious new study describes the full chain of events by which ocean biogeochemical changes triggered by manmade greenhouse gas emissions may cascade through marine habitats and organisms, penetrating to the deep ocean and eventually influencing humans.

Previous analyses have focused mainly on [ocean warming](#) and acidification, considerably underestimating the biological and social consequences of [climate change](#). Factoring in predictable synergistic changes such as the depletion of dissolved oxygen in seawater and a decline in productivity of ocean ecosystems, the new study shows that no corner of the world ocean will be untouched by climate change by 2100.

"When you look at the world ocean, there are few places that will be free of changes; most will suffer the simultaneous effects of warming, acidification, and reductions in oxygen and productivity," said lead author Camilo Mora, assistant professor at the Department of Geography in the College of Social Sciences at the University of Hawai'i at Manoa (UH Manoa). "The consequences of these co-occurring changes are massive—everything from species survival, to abundance, to range size, to body size, to species richness, to ecosystem functioning are affected by changes in ocean biogeochemistry."

The human ramifications of these changes are likely to be massive and disruptive. Food chains, fishing, and tourism could all be impacted. The study shows that some 470 to 870 million of the world's poorest people rely on the ocean for food, jobs, and revenues, and live in countries

where ocean goods and services could be compromised by multiple ocean biogeochemical changes.

Mora and Craig Smith with UH Manoa's School of Ocean and Earth Science and Technology (SOEST) worked with a 28-person international collaboration of climate modelers, biogeochemists, oceanographers, and social scientists to develop the study, which is due for publication October 15 in the scientific journal *PLOS Biology*.

The researchers used the most recent and robust models of projected climate change developed for the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC) to inform their analysis. They quantified the extent of co-occurrence of changes in temperature, pH, oxygen, and primary productivity based on two scenarios: a business-as-usual scenario wherein atmospheric CO<sub>2</sub> concentrations could reach 900 ppm by 2100, and an alternative scenario under which concentrations only reach 550 ppm by 2100 (representing a concerted, rapid CO<sub>2</sub> mitigation effort, beginning today).

They discovered that most of the world's ocean surface will be simultaneously impacted by varying intensities of ocean warming, acidification, oxygen depletion, or shortfalls in productivity. Only a very small fraction of the oceans, mostly in polar regions, will face the opposing effects of increases in oxygen or productivity, and nowhere will there be cooling or pH increase.

"Even the seemingly positive changes at high latitudes are not necessary beneficial. Invasive species have been immigrating to these areas due to changing ocean conditions and will threaten the local species and the humans who depend on them," said co-author Chih-Lin Wei, a postdoctoral fellow at Ocean Science Centre, Memorial University of Newfoundland, Canada.

The researchers assembled global distribution maps of 32 [marine habitats](#) and biodiversity hotspots to assess their potential vulnerability to the changes. As a final step, they used available data on human dependency on ocean goods and services and social adaptability to estimate the vulnerability of coastal populations to the projected ocean biogeochemical changes.

"Other studies have looked at small-scale impacts, but this is the first time that we've been able to look the entire [world ocean](#) and how co-occurring stressors will differentially impact the earth's diverse habitats and people," said co-author Andrew Thurber, a postdoctoral fellow at Oregon State University. "The real power is in the quantitative, predictive approach using IPCC climate models that allow us to see how much it will all change, and also how confident we can be in our estimates."

By 2100, global averages for the upper layer of the ocean could experience a temperature increase of 1.2 to 2.6° C, a dissolved oxygen concentration reduction of ~2% to 4% of current values, a pH decline of 0.15 to 0.31, and diminished phytoplankton production by ~4% to 10% from current values. The seafloor was projected to experience smaller changes in temperature and pH, and similar reductions in dissolved oxygen.

Of the many marine habitats analyzed in the study, researchers found that coral reefs, seagrass beds, and shallow soft-bottom benthic habitats would experience the largest absolute changes in ocean biogeochemistry, while deep-sea habitats would experience the smallest changes.

Co-author Lisa Levin, a professor at Scripps Institution of Oceanography at the University of California, San Diego, notes: "Because many deep-sea ecosystems are so stable, even small changes in temperature, [oxygen](#), and pH may lower the resilience of deep-sea communities. This is a

growing concern as humans extract more resources and create more disturbances in the deep ocean."

"The deep-sea floor covers most of the Earth's surface and provides a whole host of important ecosystem services including carbon sequestration in seafloor sediments, buffering of ocean acidity, and providing an enormous reservoir of biodiversity," said Smith.

"Nonetheless, very little attention has been paid to modeling the effects of climate change on these truly vast ecosystems. Perhaps not surprisingly, many deep seafloor ecosystems appear susceptible to the effects of climate warming over the next century."

"The impacts of climate change will be felt from the [ocean](#) surface to the seafloor. It is truly scary to consider how vast these impacts will be," said co-author Andrew K. Sweetman, who helped to convene the original team of investigators and now leads the deep-sea ecosystem research group at the International Research Institute of Stavanger, Norway. "This is one legacy that we as humans should not be allowed to ignore."

**More information:** Mora C, Wei C-L, Rollo A, Amaro T, Baco AR, et al. (2013) Biotic and Human Vulnerability to Projected Changes in Ocean Biogeochemistry over the 21st Century. *PLoS Biol* 11(10): e1001682. doi:10.1371/journal.pbio.1001682

Provided by University of Hawaii at Manoa

Citation: World ocean systems undermined by climate change by 2100 (2013, October 15)  
retrieved 10 April 2024 from  
<https://phys.org/news/2013-10-world-ocean-undermined-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private
---

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.