


# ATLAS sees Higgs boson decay to fermions

November 28 2013, by Sylvie Brunet, Abha Eli Phoboo

---



The ATLAS detector, open during a recent technical stop. Credit: Maximilien Brice/CERN


The ATLAS experiment at CERN has released preliminary results that show evidence that the Higgs boson decays to two tau particles. Taus belong to a group of subatomic particles called the fermions, which make up matter. This result – measured at 4.1 sigma on the 5-point scale particle physicists use to determine the certainty of a result – is the first

evidence for a Higgs decay to fermions.

On 4 July 2012, the ATLAS and CMS experiments at CERN announced the discovery of a new particle, which was later confirmed to be a Higgs boson.

For physicists, the discovery meant the beginning of a quest to find out what the new particle was, if it fit in the Standard Model, our current [model](#) of nature in [particle physics](#), or if its properties could point to new physics beyond that model. An important property of the Higgs boson that ATLAS physicists are trying to measure is how it decays.

The Higgs boson lives only for a short time and disintegrates into other [particles](#). The various possibilities of the final states are called decay modes. So far, ATLAS physicists had found evidence that the Higgs boson decays into different types of gauge bosons - the kind of elementary particles that carry forces. The other family of fundamental particles, the [fermions](#), make up matter. The tau is a fermion and behaves like a very massive electron.



Graphical representation of a Higgs boson decaying to two tau particles in the ATLAS detector. The taus decay into an electron (blue line) and a muon (red line) Credit: ATLAS

The Brout-Englert-Higgs mechanism was first proposed to describe how gauge bosons acquire mass. But the Standard Model predicts that fermions also acquire mass in this manner, so the Higgs boson could decay directly to either bosons or fermions. The new preliminary result from ATLAS shows clear evidence that the Higgs boson indeed does decay to fermions, consistent with the rate predicted by the Standard Model.

This important finding was made possible through careful analysis of data produced by the LHC during its first run. Only with new data will physicists be able to determine if the compatibility remains or if other new models become viable. Fortunately, the next LHC run, which begins

in 2015, is expected to produce several times the existing data sample. In addition, the proton collisions will be at higher energies, producing Higgs bosons at higher rates.

ATLAS' broad physics programme, which includes precision measurements of the Higgs boson, will continue to test the Standard Model. The years ahead will be exciting for particle physics as – the LHC experiments have found new territory that they have only just begun to explore.

**More information:** [atlas.ch/news/2013/higgs-into-fermions.html](https://atlas.ch/news/2013/higgs-into-fermions.html)

Provided by CERN

Citation: ATLAS sees Higgs boson decay to fermions (2013, November 28) retrieved 25 April 2024 from <https://phys.org/news/2013-11-atlas-higgs-boson-fermions.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--