


Brine linked to glacial health

December 6 2013, by Carol Rasmussen



This image shows the Totten Glacier ice shelf in East Antarctica (the wrinkled white area at top left) on Sept. 25, 2013. Two large open-water polynyas appear on the sea ice below and to the right of the shelf, as well as several smaller ones. The open-water areas are bright black. The stippled diagonal line from lower left to upper right is the outer edge of the sea ice, with cloud cover to the right of that line. The image is from the Moderate Resolution Imaging Spectroradiometer

instrument on NASA's Aqua satellite. Credit: NASA.

(Phys.org) —A new NASA-led study has discovered an intriguing link between sea ice conditions and the melting rate of Totten Glacier, the glacier in East Antarctica that discharges the most ice into the ocean. The discovery, involving cold, extra salty water - brine - that forms within openings in sea ice, adds to our understanding of how ice sheets interact with the ocean, and may improve our ability to forecast and prepare for future sea level rise.

"I was curious why Totten was changing so fast when the glacier just next to it wasn't changing much," said Ala Khazender of NASA's Jet Propulsion Laboratory, Pasadena, Calif., lead author of the new study, published online Dec. 5 in the journal *Nature Communications*. Combining satellite observations with ocean numerical modeling, Khazender and his colleagues developed a hypothesis that reductions in the volume of brine would increase Totten's thinning and melting. Additional research supported that hypothesis.

Ice loss seen in Antarctica is generally attributed to the well-documented rise in temperature of the surrounding ocean, but scientists are still puzzling out the mechanisms behind the regional variations that they are observing. The new study highlights the key role of processes occurring on small geographic scales in determining how global climate change can affect the stability of [ice](#) sheets.

Satellite observations from NASA's ICESat-1, which measures how much ice surfaces are rising or falling over time, revealed that Totten Glacier was thinning rapidly. It currently discharges enough ice into the surrounding ocean to fill Lake Erie in just over a week. The nearby Moscow University Glacier and its floating ice shelf were showing little

change. Why the difference? "We were convinced that the answer must be in the ocean," Khazender said.

The ocean around Antarctica is warmer than both the continent's icy surface and the polar air. Ice shelves (the floating front edges of glaciers that extend tens to hundreds of miles offshore) melt more because of contact with ocean water below them than they do because of sunlight. Melting at the undersides of [ice shelves](#) is part of Antarctica's natural water cycle, but when glaciers start melting unusually quickly, it's a sign that something is off balance.

Khazender and his team of colleagues from JPL; UCLA; the University of California, Irvine; and Utrecht University in the Netherlands combined ICESat remote sensing observations from 2003 to 2008 with ocean numerical computer models to seek insights into the interaction between the ice shelves and their ocean basin.

That ocean basin, as elsewhere around Antarctica, contains polynyas (poe-LEEN-yahs), large, annually recurring openings in the winter sea ice cover. Polynya sizes and numbers vary markedly from winter to winter, although there is no overall trend in this region. The computer simulations revealed that these year-to-year variations in the polynyas greatly affected the glacier's melting rate.

In polynyas, large quantities of sea ice form, only to be swept away by the winds that formed the openings in the first place. When seawater freezes it expels its salts, producing a layer of very dense, briny water at the freezing temperature. The cold and dense brine formed in polynyas sinks to the seafloor, where it can flow into the cavities under the ice shelves, just as warmer ocean water could.

The researchers hypothesized that when the cold brine pooled under Totten Ice Shelf, it mixed with the water there, lowering its temperature

and slowing the glacier's winter melt rate. If so, a reduction in cold brine would mean the glacier's winter melt rate would increase.

The team then examined a data set of passive microwave measurements from the Defense Meteorological Satellite Program. These showed that in the latter part of the study period, the extent of polynyas (and therefore the production of cold brine) decreased significantly. ICESat observations showed that at the same time, the thinning of Totten Glacier increased, as the team's hypothesis predicted it would.

If there are more winters with reduced polynya extents, Khazender points out, the cavity under Totten can fill with warmer ocean water rather than cold brine. "If that happens, the glacier's flow could be significantly destabilized, causing it to discharge even more ice into the ocean," he said.

"With the widespread changes seen in Antarctic [sea ice](#) conditions over the last few years, this process could be affecting other glaciers around Antarctica and the volume of ice they discharge into the [ocean](#)," he added.

More information: For more information on ICESat, visit: icesat.gsfc.nasa.gov/. For more information on the ECCO2 ocean modeling and data synthesis project, visit: ecco2.jpl.nasa.gov/

Provided by NASA

Citation: Brine linked to glacial health (2013, December 6) retrieved 9 April 2024 from <https://phys.org/news/2013-12-brine-linked-glacial-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.