

When liquids behave like solids

December 10 2013, by Lisa Zyga



(First row) SEM topography images of 1-mm drops of mercury resting on three surfaces of increasing roughness. The third surface is super-mercury-phobic. (Second row) Images of mercury drops during their detachment from these surfaces. While the drops deform considerably on the first two surfaces, on the third surface the drop deforms very little. (Third row) Force data during compression and detachment cycles show a large force hysteresis on the first two surfaces, while the hysteresis is almost absent on the third one. This behavior is similar to that of solids, but very unusual for liquids. Credit: Escobar and Castillo. ©2013 American Physical Society



(Phys.org) —When a rubber ball and a droplet of water are compressed onto a solid surface, they behave very differently. For the ball, the compression process is reversible, so the ball retains its original form when decompressed. In contrast, the compression process for the water droplet is irreversible, and the droplet's contact angle with the surface irreversibly changes because of the way the droplet interacts with the surface's chemical or physical inhomogeneities.

But now in a new study, physicists have shown that a droplet of liquid <u>mercury</u> can undergo a reversible compression process like a solid object does, as long as the surface it interacts with is "super-mercury-phobic." Such a surface is very resistant to mercury, so the mercury droplet does not spread out like a typical liquid droplet does.

The physicists, Juan V. Escobar and Rolando Castillo at the National Autonomous University of Mexico in Mexico City, have published a paper on how a liquid can behave like a solid in a recent issue of *Physical Review Letters*.

"On one hand, we have developed a novel technique with huge potential to study wetting in general and in particular the force of adhesion between a liquid and a solid, and on the other hand we helped bridge two apparently separate areas of physics (surface physics and mechanics)," Escobar told *Phys.org.* "From a more fundamental point of view, it could be said that our results show how Nature does not care about the specific source of the restoring force in a process of compression: as long as it enters the energy equation in mathematically similar form, the phenomenology will be the same.

Normally, the compression of a liquid droplet onto a <u>solid surface</u> is a hysteretic process, meaning that the droplet's contact angle depends on



its compression history. However, previous studies have predicted that the contact angle hysteresis can disappear under certain conditions, essentially making the compression process reversible.

Here, the physicists experimentally confirmed this prediction by compressing mercury droplets onto different types of surfaces. On rough surfaces, the droplets still exhibit contact angle hysteresis, but the droplets on the super-mercury-phobic surface do not. The researchers attribute the vanishing contact angle hysteresis to an extremely low surface energy between the droplet and the surface.

Along with this finding, the researchers made an unexpected discovery: even though the mercury droplet's <u>contact angle</u> hysteresis has vanished, it still has a small but measurable adhesion force, which is also independent of its compression history.

Taken together, these two observations mean that the mercury droplet behaves like an elastic solid during a repeated compressiondecompression process. The researchers explain that this behavior occurs because the dewetting of a liquid can be thought of as the equivalent of the detaching of an elastic solid. These results agree with a theoretical model in which the surface tension of a liquid is the counterpart of the restoring force of an elastic solid. The experiments could have applications for further investigation of a variety of <u>liquid</u> behavior.

"Our instrument can be used to study the dynamic wetting of a wide variety of liquids and surfaces," Escobar said. "In particular, it can be used to study the dynamic formation of the so called 'pinning points' that make the contact line stuck (or pin) on the <u>surface</u> (this study is underway). It can also be used to study the phenomenon called nonplastic ageing (this study is also underway). Also, our instrument could be used with polymers as well."



More information: Juan V. Escobar and Rolando Castillo. "Force of Adhesion Upon Loss of Contact Angle Hysteresis: When a Liquid Behaves Like a Solid." *PRL* 111, 226102 (2013). DOI: 10.1103/PhysRevLett.111.226102

© 2013 Phys.org. All rights reserved.

Citation: When liquids behave like solids (2013, December 10) retrieved 2 May 2024 from <u>https://phys.org/news/2013-12-liquids-solids.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.