


Two teams independently find that adding vibration helps couple light to graphene

December 12 2013, by Bob Yirka



Credit: J. Schiefele et al., *Phys. Rev. Lett.* (2013)

(Phys.org) —Two teams of researchers, one working in Saudi Arabia, the other in Spain, have discovered independently that adding vibrations to a graphene surface allows for more efficient conversion of photons to plasmons. In their papers, both published in the journal *Physical Review Letters*, each team describes how they found that causing a two-dimensional graphene surface to vibrate (using two different methods) led to a huge increase in light energy being coupled.

It's very common and relatively easy to use light or electronics as a means of carrying data—what's difficult is using both in the same device. The problem is getting [photons](#) converted to electrons and vice-versa. Recent research has found that using two dimensional graphene as

a means for doing so might be feasible, but up until now, scientists have only been able to achieve efficiencies of approximately 2 percent. In this new effort, both teams were able to increase the efficiency to 50 percent by causing the graphene surface to vibrate in a tunable way as it was being hit with light.

Graphene is used because of its unique honeycomb structure that allows for long duration plasmons—quasi-particles which have an oscillation property—that can be tuned to desired frequencies. Current methods of coupling that rely on shaping the graphene into ribbons have been found to be highly inefficient due to scattering at the edges (and that the approach does not allow for tuning patterns.)

To reduce scattering, both teams applied a vibrating force to the [graphene sheet](#). The Saudi team attached an actuator—the Spanish team added piezoelectric material as a base to the graphene. Both resulted in the same outcome, namely, causing the electrons in the [surface](#) of the graphene to vibrate as they were being struck by photons—tuning the vibrations allowed for rousing the conduction electrons into plasmons, which thereafter could be processed by electronic components.

The results by both teams are just the first step in creating mechanisms that are able to join electronic functions with photons which may one day lead to devices such as exotic super-sensitive chemical detectors, new kinds of photovoltaic cells or general purpose nano-optoelectronics devices.

More information: Exciting Graphene Surface Plasmon Polaritons through Light and Sound Interplay, *Phys. Rev. Lett.* 111, 237404 (2013) prl.aps.org/abstract/PRL/v111/i23/e237404

Coupling Light into Graphene Plasmons through Surface Acoustic Waves, *Phys. Rev. Lett.* 111, 237405 (2013)

prl.aps.org/abstract/PRL/v111/i23/e237405

© 2013 Phys.org

Citation: Two teams independently find that adding vibration helps couple light to graphene (2013, December 12) retrieved 27 April 2024 from <https://phys.org/news/2013-12-teams-independently-adding-vibration-couple.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.