

# Effective control of invasive weeds can help attempts at reforestation in Panama

January 28 2014

---



This picture shows the ease by which the seeds of *Saccharum spontaneum* can germinate in the environment. As the seed is wind borne it can travel large distances (kms) and potentially spread the invasion. Credit: Graham D. Bonnett

*Saccharum spontaneum* is an invasive grass that has spread extensively in disturbed areas throughout the Panama Canal watershed, where it has created a fire hazard and inhibited reforestation efforts. The weed originally believed to be originally from India, is perfectly adapted to the conditions in Panama and produces excessive amounts of biomass during

the wet season, which impedes reforestation efforts. A new study published in the open access journal [NeoBiota](#) proposes an effective method for controlling the growth, based on analysis of its reproductive biology.

Currently physical removal of [aboveground biomass](#) is the primary means of controlling the weed, which is largely ineffective and does little to inhibit spread of the species. This is due to the insufficient knowledge about reproduction of the species and this is where science comes to the rescue.

A team of scientists from Australia and Panama provide a detailed examination of a series of studies looking at some of the basic reproductive mechanisms and strategies utilised by *S. spontaneum* to provide information to support development of better targeted management strategies.

It turns out that *S. spontaneum* has a very good survival toolkit being able to reproduce through buds on stems that had been dried for up to six weeks. Separate experiments showed that even leftover stem fragments could sprout when left on the surface or buried shallowly and that larger pieces sprouted more readily than smaller pieces.



This picture shows *Saccharum spontaneum* along a road close to Miraflores Locks on the Panama Canal with the cruise ship Celebrity infinity passing through the canal. Credit: Graham D. Bonnett



This picture shows the potential for spread via vegetative pieces of the plant. The plant is pictured in a pile of earth about to be moved from site to another. Credit: Graham D. Bonnett

The study shows that the through better knowledge the panacea of a big problem can turn out to be very simple. A good timing of management actions to prevent flowering would significantly reduce the seed load into the environment and help to prevent spread to new sites. Similarly simple but effective would be cutting stems into smaller pieces allowing them to dry out and reduce the ability of buds to sprout.

**More information:** Bonnett GD, Kushner JNS, Saltonstall K (2014) The reproductive biology of *Saccharum spontaneum* L.: implications for management of this invasive weed in Panama. *NeoBiota* 20: 61. [DOI: 10.3897/neobiota.20.6163](https://doi.org/10.3897/neobiota.20.6163)

Provided by Pensoft Publishers

Citation: Effective control of invasive weeds can help attempts at reforestation in Panama (2014, January 28) retrieved 11 May 2024 from <https://phys.org/news/2014-01-effective-invasive-weeds-reforestation-panama.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.