

Highly efficient broadband terahertz radiation from metamaterials

January 17 2014, by Breehan Gerleman Lucchesi



A THz spectrometer driven by femtosecond laser pulses was used to demonstrate THz emission from a split-ring resonator metamaterial of single nanometer thickness.

(Phys.org) —Scientists at the U.S. Department of Energy's Ames Laboratory have demonstrated broadband terahertz (THz) wave generation using metamaterials. The discovery may help develop noninvasive imaging and sensing, and make possible THz-speed information communication, processing and storage. The results appeared in the Jan. 8 issue of *Nature Communications*.

Terahertz electromagnetic <u>waves</u> occupy a middle ground between



electronics waves, like microwave and radio waves, and photonics waves, such as infrared and UV waves. Potentially, THz waves may accelerate telecom technologies and break new ground in understanding the fundamental properties of photonics. Challenges related to efficiently generating and detecting THz waves has primarily limited their use.

Traditional methods seek to either compress oscillating waves from the electronic range or stretch waves from the optical range. But when compressing waves, the THz frequency becomes too high to be generated and detected by conventional electronic devices. So, this approach normally requires either a large-scale electron accelerator facility or highly electrically-biased photoconductive antennas that produce only a narrow range of waves.

To stretch optical waves, most techniques include mixing two laser frequencies inside an inorganic or organic crystal. However, the natural properties of these crystals result in low efficiency.

So, to address these challenges, the Ames Laboratory team looked outside natural materials for a possible solution. They used man-made materials called metamaterials, which exhibit optical and magnetic properties not found in nature.

Costas Soukoulis, an Ames Laboratory physicist and expert in designing <u>metamaterials</u>, along with collaborators at Karlsruhe Institute of Technology in Germany, created a metamaterial made up of a special type of meta-atom called split-ring resonators. Split-ring resonators, because of their u-shaped design, display a strong magnetic response to any desired frequency waves in the THz to infrared spectrum.





A team led by Ames Laboratory physicists demonstrated broadband, gapless terahertz emission (red line) from split-ring resonator metamaterials (background) in the telecomm wavelength. The THz emission spectra exhibit significant enhancement at magnetic-dipole resonance of the metamaterials emitter (shown in inset image). This approach has potential to generate gapless spectrum covering the entire THz band, which is key to developing practical THz technologies and to exploring fundamental understanding of optics.

Ames Laboratory physicist Jigang Wang, who specializes in ultra-fast laser spectroscopy, designed the femto-second laser experiment to demonstrate THz emission from the metamaterial of a single nanometer thickness.

"The combination of ultra-short laser pulses with the unique and unusual properties of the metamaterial generates efficient and broadband THz waves from emitters of significantly reduced thickness," says Wang, who



is also an associate professor of Physics and Astronomy at Iowa State University.

The team demonstrated their technique using the wavelength used by telecommunications (1.5 microns), but Wang says that the THz generation can be tailored simply by tuning the size of the meta-atoms in the metamaterial.

"In principle, we can expand this technique to cover the entire THz range," said Soukoulis, who is also a Distinguished Professor of physics and astronomy at Iowa State University.

What's more, the team's metamaterial THz emitter measured only 40 nanometers and performed as well as traditional emitters that are thousands of times thicker.

"Our approach provides a potential solution to bridge the 'THz technology gap' by solving the four key challenges in the THz emitter technology: efficiency; broadband spectrum; compact size; and tunability," said Wang.

Soukoulis, Wang, Liang Luo and Thomas Koschny's work at Ames Laboratory was supported by the U.S. Department of Energy's Office of Science. Wang's work is partially supported by Ames Laboratory's Laboratory Directed Research and Development (LDRD) funding.

DOE's Office of Science is the single largest supporter of basic research in the physical sciences in the United States, and is working to address some of the most pressing challenges of our time. For more information, please visit the Office of Science website at science.energy.gov/.

Ames Laboratory is a U.S. Department of Energy Office of Science national laboratory operated by Iowa State University. Ames Laboratory



creates innovative materials, technologies and energy solutions. We use our expertise, unique capabilities and interdisciplinary collaborations to solve global problems.

More information: "Broadband terahertz generation from metamaterials." Liang Luo, Ioannis Chatzakis, Jigang Wang, Fabian B. P. Niesler, Martin Wegener, Thomas Koschny, Costas M. Soukoulis. *Nature Communications* 5, Article number: 3055 <u>DOI:</u> <u>10.1038/ncomms4055</u>. Received 05 August 2013 Accepted 03 December 2013 Published 08 January 2014

Provided by US Department of Energy

Citation: Highly efficient broadband terahertz radiation from metamaterials (2014, January 17) retrieved 4 May 2024 from https://phys.org/news/2014-01-highly-efficient-broadband-terahertz-metamaterials.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.