

# Mystery of California's killer rattlesnakes solved

January 28 2014

---



Southern Pacific Rattlesnake. Credit: Chip Cochran.

(Phys.org) —A surge in snakebite deaths caused by one of North America's most dangerous snakes has been baffling doctors, but new research may hold the key to saving lives.

A study of Southern Pacific Rattlesnake [venom](#), led by The University of Queensland's Associate Professor Bryan Fry, has found that by knowing the location where a person was bitten, doctors are better equipped to offer life-saving treatment.

"If clinicians know where a person was bitten, they will know how the patient is likely to be affected," Dr Fry said.

"These snakes live in habitats as diverse as the isolated Catalina Island, the high-altitude San Jacinto mountains, the grassy hills of Loma Linda and the desert transition zone of Phelan.

"In a two-hour drive from the desert floor to the top of the San Jacinto Mountains, the venom goes from destroying the blood to frying the nerves instead.

"Over millions of years, living in these very different habitats has led to specialised venom chemistry which needs to be understood to effectively treat snakebite patients."

Dr Fry and his team sampled venoms from four diverse regions, including Catalina Island, and analysed the venom chemistry and evolution.

The team found significant differences between populations.

"Mapping the geographical venom variation of this species has important implications for treating bites," he said.

Dr Fry said the study disproved previous reports that recent, rapid change in the venom was the cause, leading to increased reports of unusual and highly toxic effects in patients.

"Clinicians had been at a loss to explain what was happening," Professor Fry said.

"However, rather than changing rapidly, the venom varies dramatically between different populations of the snakes due to long-term adaptation

to different environments."

Dr Fry also said reports of unusual effects were likely due to better record keeping and reporting and changes in human behavior, rather than any recent changes to the venom.

"New housing estates are being built on what used to be remote areas, and people with low snake awareness are coming into close contact with Southern Pacific Rattlesnakes," he said.

"Many times when a patient presents at the hospital it is because they tried to kill the snake and were bitten in the process."

**More information:** Kartik Sunagar, Eivind A.B. Undheim, Holger Scheib, Eric C.K. Gren, Chip Cochran, Carl E. Person, Ivan Koludarov, Wayne Kelln, William K. Hayes, Glenn F. King, Agosthino Antunes, Bryan Grieg Fry. "Intraspecific venom variation in the medically significant Southern Pacific Rattlesnake (*Crotalus oreganus helleri*): Biodiscovery, clinical and evolutionary implications." *Journal of Proteomics*, Available online 24 January 2014, ISSN 1874-3919, [dx.doi.org/10.1016/j.jprot.2014.01.013](https://doi.org/10.1016/j.jprot.2014.01.013).

Provided by University of Queensland

Citation: Mystery of California's killer rattlesnakes solved (2014, January 28) retrieved 19 April 2024 from <https://phys.org/news/2014-01-mystery-california-killer-rattlesnakes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.