

Using rare earths to interpret certain fossils

January 30 2014

Until now, interpreting flattened fossils was a major challenge. Now, a new approach for the analysis of such fossils has been developed by a team bringing together researchers from the IPANEMA unit (CNRS), the Centre de Recherche sur la Paléobiodiversité et les Paléoenvironnements (CNRS) and the SOLEIL synchrotron. This non-destructive method makes use of chemical elements known as rare earths. By locating and quantifying such elements in trace amounts, it is possible to improve interpretation of fossil morphology. This enabled the researchers to describe not only the anatomy but also the environment of preservation of three fossils of Cretaceous age.

Published on 29 January in the journal *Plos One*, the work should facilitate the interpretation of many flattened fossils, especially those that are exceptionally well conserved.

During the [fossilization](#) process, animal and plant remains are frequently flattened and compressed into two dimensions by the pressure of rocks, which sometimes proves to be a real obstacle to the study of such fossils. An added difficulty is that such flattened fossils undergo physical and chemical alteration during fossilization, making it even harder to interpret them. However, such fossils can contain invaluable information. In particular, when their anatomy is well preserved (in which case they are termed exceptionally well-preserved fossils), soft tissues such as muscles are also fossilized. However, locating such tissues remains particularly difficult due to the limited contrast attained in optical microscopy and to the limits of tomography, techniques that are today commonly used to study fossils.

Now, researchers from CNRS, the French Natural History Museum (MNHN) and the SOLEIL synchrotron have designed and developed a novel non-destructive approach based on the localization of [rare earths](#). These chemical elements (yttrium and the lanthanides) are known to exist in fossils in trace amounts, typically from 1 to 1000 micrograms per gram of material. However, depending on the type of tissue, the quantity of trace elements incorporated during fossilization varies. Such preferential fixing makes it possible to distinguish between the anatomical parts of a fossil. This appears as a strong contrast between different [chemical elements](#) according to the types of fossil tissue, when the fossil is characterized using synchrotron fast X-ray fluorescence imaging. To speed up the analysis, the team proposes a rapid method for differentiating tissue, based on the probabilistic nature of the data measured.

The scientists applied this approach to three fossils (two fishes and a shrimp) discovered in Morocco and dating from the Upper Cretaceous, around 100 million years ago. The contrasts revealed using this method enabled them to distinguish between hard tissues (bones and shells) and [soft tissues](#) (muscles and other fossilized organs). In particular, this enabled them to discover the previously hidden anatomical features of a fossil fish of which only one specimen is known, one of whose skull bones is in the form of a wide, notched blade.

This new approach makes it possible to obtain a detailed, accurate view of the anatomy of a fossil without damaging it and without the need for prior delicate sample preparation. It is particularly suitable for flattened fossils given that X-rays penetrate the fossil to a depth of a few fractions of a millimeter. The technique also revealed certain bones concealed under a fine layer of rock, enabling them to be viewed directly. For instance, it enabled certain concealed appendages of a fossil shrimp to be viewed, such as the legs and the antennae, which hold important information for the study of its relationship to other shrimps. In addition,

rare earth content reflects the environment in which a fossil is preserved, such as connectivity to surrounding water networks, local physico-chemical conditions, and the properties of the mineral phases making up the fossils, thus enabling them to be better described.

The work should therefore facilitate the interpretation of flattened fossils, which are very common in the fossil record. It opens up new prospects not only for paleoenvironmental studies but also for a better understanding of long-term fossilization processes.

More information: "Trace elemental imaging of rare earth elements discriminates tissues at microscale in flat fossils," Pierre Gueriau, Cristian Mocuta, Didier B. Dutheil, Serge X. Cohen, Dominique Thiaudière, The OT1 consortium, Sylvain Charbonnier, Gaël Clément & Loïc Bertrand. *PLoS ONE*, 29 January 2014.

Provided by CNRS

Citation: Using rare earths to interpret certain fossils (2014, January 30) retrieved 26 June 2024 from <https://phys.org/news/2014-01-rare-earths-fossils.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.