

Scientists improve detection of Arctic clouds

January 7 2014



Caught in the early morning light, the multifilter rotating shadowband radiometer instrument takes measurements of direct and diffuse sunlight in Barrow, Alaska. Now that PNNL scientists have devised a new method for screening out the thin clouds prevalent at this Alaska site, those measurements will be much more effective. Credit: ARM Climate Research Facility.

(Phys.org) —Thin Arctic clouds can no longer hide, thanks to scientists at Pacific Northwest National Laboratory. Atmospheric data gathered by skyward pointing instruments can be "contaminated" by clouds so wispy

that they appear to the instruments as tiny, suspended particles called aerosols, the desired target. The researchers showed how a simple modification of typical methods used to detect denser clouds and remove them from the data results in more accurate measurements of aerosols. (See sidebar, Cloud Contamination and Cloud Screening). With these improvements, scientists can better understand how aerosols influence climate changes in the Arctic.

"Thin and almost uniform Arctic clouds are occasionally undetected by well-established methods," said Dr. Evgueni Kassianov, atmospheric scientist and lead author of the study. "Our research found a minor modification of these methods that substantially improves the detection of these clouds, thereby reducing cloud contamination of aerosol data sets."

Sunbathers see clouds as an obstacle between them and the sun's bronzing rays. Scientists have a similar point of view when trying to capture atmospheric information from the ground by using instruments that point up. In some cases, the upward looking instruments require clear skies to gather the correct data. Clouds can hinder this requirement by blocking the instruments' view. To make matters worse, if the clouds are thin and wispy, these clouds may contaminate observations thought to have been made under clear sky conditions. In this study, the scientists were particularly interested in gaining information in the Arctic where thin clouds abound. The improved method proposed by these scientists will increase effectiveness of atmospheric measurements to understand possible abrupt climate changes in the Arctic.

Researchers at PNNL conducted an observational study based on a multi-year and integrated dataset of aerosol and clouds collected at the high-latitude Atmospheric Radiation Measurement (ARM) Climate Research Facility's North Slope of Alaska (NSA). Two sites provided data, one on the northern coast in Barrow, Alaska, and the second in Atkasuk, Alaska,

about 100 kilometers inland from the coast. They documented occasional failures of the well-established cloud-screening methods, which had great success at other lower-latitude ARM facility sites (where the [clouds](#) are thicker and more opaque). The team uncovered the main reasons for such failures. They proposed and evaluated an improved version of these cloud-screening methods, which in turn led to improved aerosol climatology at the ARM Alaska sites.

Researchers plan to apply the improved method to similar datasets collected at ARM sites globally. These extended applications will provide important observational constraints for model-based estimates of the aerosol impact on the Earth's radiation budget.

More information: Kassianov E, C Flynn, A Koontz, C Sivaraman, and J Barnard. 2013. "Failure and Redemption of Multifilter Rotating Shadowband Radiometer (MFRSR)/Normal Incidence Multifilter Radiometer (NIMFR) Cloud Screening: Contrasting Algorithm Performance at Atmospheric Radiation Measurement (ARM) North Slope of Alaska (NSA) and Southern Great Plains (SGP) Sites." *Atmosphere* 4(3):299-314. [DOI: 10.3390/atmos4030299](https://doi.org/10.3390/atmos4030299)

Provided by Pacific Northwest National Laboratory

Citation: Scientists improve detection of Arctic clouds (2014, January 7) retrieved 12 May 2024 from <https://phys.org/news/2014-01-scientists-arctic-clouds.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
