

## Soil microbes alter DNA in response to warming

January 16 2014



Soil plots that were heated to simulate the rise in temperatures forecast by climate models. The radiators (top) warm the plots 2 degrees Celsius. Credit: Mengting Yuan, U. of Oklahoma

As scientists forecast the impacts of climate change, one missing piece of the puzzle is what will happen to the carbon in the soil and the microbes that control the fate of this carbon as the planet warms.

Scientists studying grasslands in Oklahoma have discovered that an



increase of 2 degrees Celsius in the air temperature above the soil creates significant changes to the microbial ecosystem underground. Compared to a control group with no warming, plants in the warmer plots grew faster and higher, which put more <u>carbon</u> into the soil as the plants senesce. The microbial ecosystem responded by altering its DNA to enhance the ability to handle the excess carbon.

"What we conclude from this study is the warming has an effect on the soil ecosystem," said Kostas Konstantinidis, an assistant professor who holds the Carlton S. Wilder Chair in Environmental Engineering at the Georgia Institute of Technology. "It does appear that the microbes change genetically to take advantage of the opportunity given to them."

The study was published online Dec. 27, 2013, in the journal *Applied and Environmental Microbiology*. The research was sponsored by the Department of Energy, and involved collaboration with several universities, including the University of Oklahoma.

The findings are the culmination of a 10-year study that seeks to understand how the most intricate ecosystem in nature—soil—will respond to <u>climate change</u>.

A single gram of soil is home to a billion bacterial cells, representing at least 4,000 different species. In comparison, the human gut is home to at least 10 times fewer different species of bacteria. Scientists have little idea what microbes in the soil do, how they do it, or how they respond to changes in their environment, Konstantinidis said. This limits the predictive capabilities of climate models.

"In models of climate change it is a black box what happens to the carbon in soil," Konstantinidis said. "One reasons that models of climate change have such big room for variation is because we don't understand the microbial activities that control carbon in the soil."





Georgia Tech assistant professor Kostas Konstantinidis (left) and former graduate student and study lead author Chengwei Luo examine *E. coli* cells. Credit: Gary Meek

Complicating matters, 99.9 percent of the microbes in the soil cannot be grown in the lab, so scientists must study them where they live. The molecular and genomic techniques to do so are a specialty of the



Konstantinidis lab.

The researchers traveled to the Kessler Farm Field Laboratory in McClain County, Oklahoma, where they conducted their study on grassland soils, which had been abandoned for agriculture use for more than 20 years. The scientists warmed plots of soil with radiators set a few feet above the ground for 10 consecutive years. They warmed these plots 2 degrees Celsius, which many <u>climate models</u> forecast as the global temperature increase over the next 50 years.

The researchers took samples of the plants, measured the carbon content and the number of microbes in the soil, and documented any changes in the warm plots versus the control plots. The team also extracted DNA from the soil and identified the genetic composition and changes of the microbes living there.

The plants in the warm plots grew better and higher. As the plants started senescing at the end of the season, their higher biomass led to more carbon in the soil. However, the <u>microbial communities</u> had increased their rate of respiration, which converted soil organic carbon to carbon dioxide (CO2), so the total carbon in the warm and control soils was similar.

The microbial communities in the warm soils had undergone significant changes during the decade of the experiment, which facilitated their higher respiration rate. For instance, the study of the DNA of the microbes revealed found that the microbial communities of the warm plots had more genes related to carbon respiration than the microbes in the control plots.

"That was consistent with the idea that the additional carbon from the plants was all respired and converted to CO2," Konstantinidis said. "We saw that the warmed microbial community was more efficient in eating



up the plant-derived soil carbon and making it CO2."

The research team plans to do similar studies in other agricultural soils and in colder areas, such as Alaska tundra permafrost ecosystems, where there is more organic carbon in the <u>soil</u>.

"There are complex interactions between plants and <u>microbes</u> and we need to understand them better to have a more predictive understanding of what's going on," Konstantinidis said. "This is the first study trying to do that, but we are not close to the complete understanding yet."

**More information:** Chengwei Luo, et al., "Soil microbial community responses to a decade of warming as revealed by comparative metagenomics," *Applied and Environmental Microbiology*, January 2013. dx.doi.org/10.1128/AEM.03712-13

Provided by Georgia Institute of Technology

Citation: Soil microbes alter DNA in response to warming (2014, January 16) retrieved 2 May 2024 from <u>https://phys.org/news/2014-01-soil-microbes-dna-response.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.