

Drastic chemical change occurring in birth of planetary system: Has the solar system also experienced it?

February 13 2014



Illustration of rotating-infalling gas toward a protostar. The abundance of sulfur monoxide is enhanced at the outer edge (colored blue) of a protoplanetary disk. Credit: The University of Tokyo

A new star is formed by gravitational contraction of an interstellar molecular cloud consisting of gas and dust. In the course of this process, a gas disk (protoplanetary disk), whose size is on the order of 100 AU, forms around the protostar and evolves into a planetary system. The solar system was also formed in this way about 4.6 billion years ago, and life was eventually born on the Earth. How unique in the universe is the situation which happened for the solar system?

In order to answer this question, understanding the formation of protoplanetary disks as well as the associated chemical evolution in various star forming regions is essential. There have been extensive observational efforts made toward this goal. So far, most of them have focused on changes in the physical structure and the kinematics during the formation process. However, it was very difficult to distinguish the [protoplanetary disk](#) from the infalling envelope clearly with such conventional approaches.

On the other hand, the chemical evolution associated with disk formation has scarcely been studied observationally because of the insufficient sensitivity and spatial resolution of previous radio telescopes. As a result, a chemical model calculation with many assumptions is the only approach. Naturally, the physical and chemical changes in the disk formation should be coupled with each other. The disk formation around a young protostar has been explored from a novel point of view looking at physics and chemistry simultaneously.

L1527 in the Taurus [molecular cloud](#) is a molecular cloud core which harbors a young protostar. A global team led by Dr. Nami Sakai, the University of Tokyo, conducted high-sensitivity, high-spatial-resolution observations of L1527 with ALMA (Atacama Large Millimeter/submillimeter Array) newly constructed in the Atacama desert in Chile, and investigated the disk formation process by using the spectral lines of several molecules.

As a result, Sakai al. have found that carbon-chain molecules and their related species such as cyclic-C₅H₂ almost completely disappear from the gas phase inside a radius of 100 AU around the protostar (Figure 1, top left; top right).




Figure 1. An infrared image of the protostar L1527 taken by the Spitzer Space Telescope. Credit: J. Tobin/NASA/JPL-Caltech

Precise measurements of the motion of the gas using the Doppler shift in

the spectral lines of the gas components revealed that 100 AU corresponds to the radius of the centrifugal barrier (Figure 2).




Figure 2. L1527 observed by Spitzer (Left) and the distributions of cyclic-C₃H₂ (center) and SO (right) observed by ALMA. ALMA reveals the gas distribution just close to the protostar. Emission from cyclic-C₃H₂ is weak toward the protostar but strong at the northern and southern parts. Meanwhile, SO has its emission peak near the protostar. Credit: J. Tobin/NASA/JPL-Caltech, N. Sakai/The University of Tokyo

At this radius, infalling gas is stopped and accumulated due to the centrifugal force, and then is gradually transferred to the inner disk. Namely, this is the edge of the disk forming region. It has clearly been indentified with the spectral line of cycle-C₅H₂.

On the other hand, the distribution of sulfur monoxide molecules (SO) is found to be localized in a ring structure located at the radius of the centrifugal barrier (100 AU) (Figure 1, bottom left; bottom right). Furthermore, the temperature of the SO molecules is found to be higher than that of the infalling gas. This means that the infalling gas probably causes a weak shock when in rushes into the outer edge of the disk at the centrifugal barrier. The gas temperature is raised around this radius, and the SO molecules frozen on dust grains are liberated into the gas phase.

Hence, the spectral line of SO also highlights the disk-formation front. Since the density of the disk is 10^8 cm^{-3} or higher, most of the molecules are frozen out onto dust grains in the disk after they pass through the front.

It is not at all anticipated that such a drastic chemical change occurs in the transition zone between the infalling envelope and the inner disk. The disk formation and the associated chemical change have successfully been detected by observations of the two chemical species, cyclic-C₅H₂ and SO.

This study has demonstrated a drastic change in chemical composition associated with disk formation around the young protostar (cf; Figure 3). With a coupled view of physics and chemistry, it has also succeeded in highlighting the outermost part of the disk where the [gas](#) is still accreting. This success was realized by high-sensitivity and high-spatial-resolution observations with ALMA, and such a study will be extended to other various star-forming regions. In particular, it is very interesting to examine how widely applicable the picture seen in L1527 is to other star-forming regions.

Although many observational efforts aimed at understanding planetary-system formation have been made, this study is novel in focusing on the chemical change. By extending this new method to various solar-type protostars using ALMA, the diversity and generality of the chemical evolution from interstellar matter to planetary matter will be unveiled within the next few years. Then, we can critically examine whether the solar system experienced this drastic chemical change. In parallel to the astronomical approach, the origin of the solar system is being investigated by exploring the [solar system](#) itself through microanalyses of meteorites, spectroscopy of comets, sample return missions to the asteroids, and so on. The present study will also have a strong impact on these studies by tracing the origins back to interstellar clouds.

More information: Nami Sakai, Takeshi Sakai, Tomoya Hirota, Yoshimasa Watanabe, Cecilia Ceccarelli, Claudine Kahane, Sandrine Bottinelli, Emmanuel Caux, Karine Demyk, Charlotte Vastel, Audrey Coutens, Vianney Taquet, Nagayoshi Ohashi, Shigehisa Takakuwa, Hsi-Wei Yen, Yuri Aikawa & Satoshi Yamamoto, "Change in the chemical composition of infalling gas forming a disk around a protostar", *Nature*, DOI: [10.1038/nature13000](https://doi.org/10.1038/nature13000).

Provided by National Astronomical Observatory of Japan

Citation: Drastic chemical change occurring in birth of planetary system: Has the solar system also experienced it? (2014, February 13) retrieved 26 April 2024 from <https://phys.org/news/2014-02-drastic-chemical-birth-planetary-solar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.