

## Better diagnosis of parasitic infections via ultrasound enrichment of rare cells

March 28 2014



(Phys.org) —Parasitic infections like malaria and sleeping sickness affect hundreds of millions of people, primarily in the poorest regions of the world. Diagnosis of these diseases is often difficult because the concentration of parasites in the blood can be very low. British scientists have now developed a simple chip-based method for enriching rare cells in blood samples. As they report in the journal *Angewandte Chemie*, this allows the detection limit for the parasites that cause malaria and sleeping sickness to be lowered by two to three orders of magnitude.



Existing techniques for the separation and enrichment of parasites in blood samples are difficult to use in isolated regions and developing countries because they usually require complex chemistry for labeling cells, costly instruments, or extensive infrastructure. An inexpensive technique that requires only small amounts of power, works without labeling the cells, and uses just a drop of blood from a fingertip, is needed. A team headed by Jonathan M. Cooper at the University of Glasgow has now developed such an approach. Their innovative method is based on an acoustically controlled microchip that is used in a battery-driven, hand-held device. The researchers successfully used their technique to enrich malaria-infected <u>blood cells</u> and the parasite that causes sleeping sickness in <u>blood samples</u>.

The chip contains a special electrode that produces ultrasound when a voltage is applied. If a drop of liquid is placed in a specific location on the device, the form of the acoustic field elicits a particular pattern of flow within the drop: a circular rotational motion. Particles whose density is lower than that of the liquid are carried against gravity with the upward rising current and transported toward the outer edge of the drop, where they accumulate. In contrast, particles with a higher density collect in the center of the droplet, because they cannot be lifted up.

This works for cells too. Red blood cells infected with the malaria parasite are less dense than non-infected cells. If the density of the drop of blood being examined is adjusted by simply adding a small amount of reagent, the acoustic chip allows the infected red blood cells to be concentrated by a factor of one hundred to one thousand at the outer edge of the blood drop. The non-infected <u>red blood cells</u> remain at the center of the drop. The method is also suitable for concentrating freeswimming parasites in blood. The researchers were able to enrich trypanosomes, the pathogens that cause <u>sleeping sickness</u>, by using their acoustic chip. Simple staining techniques then make it possible to detect the parasites.



In the future, the technique may be adapted to allow other infectious diseases and rare circulating tumor cells to be detected more readily use of this new technology.

**More information:** Bourquin, Y., Syed, A., Reboud, J., Ranford-Cartwright, L. C., Barrett, M. P. and Cooper, J. M. (2014), "Rare-Cell Enrichment by a Rapid, Label-Free, Ultrasonic Isopycnic Technique for Medical Diagnostics." *Angew. Chem. Int. Ed.* DOI: 10.1002/anie.201310401

Provided by Angewandte Chemie

Citation: Better diagnosis of parasitic infections via ultrasound enrichment of rare cells (2014, March 28) retrieved 23 June 2024 from <u>https://phys.org/news/2014-03-diagnosis-parasitic-infections-ultrasound-enrichment.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.