


Earliest evidence of limb bone marrow in the fin of a 370 million year old fish

March 19 2014



Credit: © Sophie Sanchez

This week in the journal *Proceedings of the Royal Society B*, a team of French and Swedish researchers present the earliest fossil evidence for the presence of bone marrow in the fin of a 370 million-year-old fish.

Long bones, which are found in the limb of tetrapods, are not only important for locomotion and supporting the weight of the body, but also host the bone marrow. The latter plays a major role in haematopoiesis, i.e. the formation of blood cells. In a healthy adult human, about a hundred billion to one trillion new blood cells are produced every day to maintain the stable blood circulation. The bone marrow also has an important role as part of the immune system. Although long bones are a rich source for [marrow transplantation](#), the establishment the bone marrow in the medullary cavity and its interactions with the surrounding bone still remain partly mysterious and its evolution is not well understood.

Researchers from Uppsala University in Sweden and the European Synchrotron Radiation Facility (ESRF) in France decided to look for the origin of the bone marrow within vertebrates, using synchrotron microtomography to investigate the interior structure of fossil long bones without damaging them. They discovered that *Eusthenopteron*, a Devonian (370 million year old) lobe-finned fish from Miguasha in Canada that is closely related to the first tetrapods, already exhibited typical marrow processes inside its humerus (upper arm bone). These processes are longitudinal, larger than blood vessel canals, and connect to the shoulder and elbow joint surfaces of the humerus. Thanks to the beam power of the ESRF, they were able to reach submicron resolutions and accurately reconstruct the 3D arrangement of the long-bone microanatomy of this close relative of tetrapods.

"We have discovered that the bone marrow certainly played a major role in the elongation of fin bone through complex interactions with the [trabecular bone](#)" says Sophie Sanchez, a researcher from Uppsala

University and the ESRF. "This intimate relationship, which has been demonstrated by molecular experiments in extant mammals, is actually primitive for tetrapods".

This discovery is very important for understanding the evolutionary steps that built up the distinctive architecture of tetrapod limb bones and created a location for the distinctive, complex and functionally important tissue that is [bone marrow](#). It is also a powerful demonstration of the capabilities of synchrotron microtomography.

"Without the 3D information provided by the synchrotron, we could never have understood the internal organization of the marrow space" says Per Ahlberg from Uppsala University. "If you cut a slice through a bone like this, which would damage it irreparably, you would only see an uninformative pattern of holes in the cut surface. With the synchrotron we can image the whole internal structure and understand how the marrow processes are organized, without doing any damage to the bone at all."

More information: Sanchez S, Tafforeau P, Ahlberg PE. (2014) The humerus of Eusthenopteron: a puzzling organization presaging the establishment of tetrapod limb bone marrow. Proc. R. Soc. B 281: 20140299. [dx.doi.org/10.1098/rspb.2014.0299](https://doi.org/10.1098/rspb.2014.0299)

Provided by Uppsala University

Citation: Earliest evidence of limb bone marrow in the fin of a 370 million year old fish (2014, March 19) retrieved 19 April 2024 from <https://phys.org/news/2014-03-earliest-evidence-limb-bone-marrow.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.