

Faster screening test to identify tuberculosis

March 3 2014, by Shawn Eyre




Figure. Nanotrap. (a) The antigens, CFP-10, secreted by *M. tuberculosis* were isolated by the well-engineered nanoporous thin film (b) The abundant proteins (green) in complex biological fluids were washed away. CFP-10 (orange) left inside the nanopores were cleaved by enzyme and then detected with mass spectrometry.

With 9 million new cases and 2 million deaths annually, Tuberculosis is the second most prevalent and deadliest infectious disease worldwide. As an airborne disease, it spreads easily and is very contagious. Quick detection and identification is the key to success in preventing the spread of the disease.

Conventional tuberculosis screenings suffer from low sensitivity, specificity, and high-cost. The gold standard for the diagnosis of [active tuberculosis](#) still relies on time-consuming culture tests that take 10 to 40 days to complete the screening process.

Recent nanopore research by Assistant Professor Hung-Jen Wu of the Artie McFerrin Department of Chemical Engineering at Texas A&M University, along with the Houston Methodist Research Institute, has led to a new screening tool that quickly identifies the M. tuberculosis antigen, CFP-10.

"CFP-10 is an antigen secreted by the M. tuberculosis bacteria," says Wu. "Its rod shape allows us to use various types of nanopore thin films to very easily and quickly, isolate and identify the strand using a mass spectrometer."

Wu has engineered the nanopore size, structure, and surface properties to isolate up to 90% of CFP-10 in biological samples. The whole screening process takes about nine hours including incubation time. This overall approach offers hope not only for speeding up diagnosis of active tuberculosis but also for future screening of other [infectious diseases](#).

Provided by Texas A&M University

Citation: Faster screening test to identify tuberculosis (2014, March 3) retrieved 4 May 2024 from <https://phys.org/news/2014-03-faster-screening-tuberculosis.html>

| |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|