

## **Geographers create 'easy button' to calculate river flows from space**

March 18 2014, by Meg Sullivan



Scientists can measure the flow of this Greenland river using only satellite imagery.

(Phys.org) —The frustrated attempts of a UCLA graduate student to quantify the amount of water draining from Greenland's melting ice sheet led him to devise a new way to measure river flows from outer space, he and his professor report in a new study.



The new approach relies exclusively on the measurements of a river's width over time, which can be obtained from freely available <u>satellite</u> <u>imagery</u>.

Currently, hydrologists calculate a river's discharge—the volume of water running through it at any given time—by taking a series of measurements on the ground, including not just the river's width but also its depth at multiple points and the velocity of its flows. Politics, financial considerations and terrain often stand in the way of obtaining such measurements.

"Our new method doesn't require access to the country or getting in the river to safely take measurements in the field," said Colin Gleason, lead author of the study and a doctoral student in geography in UCLA's College of Letters and Science. "As long as we can get multiple pictures of a river and apply this method to them, we can tell you how much water was flowing in the river at the time the images were taken."

The discovery is highlighted March 17 in the online edition of *Proceedings of the National Academy of Sciences (PNAS)*.

With the potential to be applied anywhere in the world, the new approach is expected to provide information that will benefit agriculture, sanitation and flood preparation, especially in countries without the resources to physically measure and monitor their <u>rivers</u>.

Additionally, it holds promise for improving the accuracy of climate models, which track recycling of water between the atmosphere and land, and ensuring that countries with trans-boundary rivers comply with international water-sharing agreements.

"This work represents a breakthrough in our ability to address one of the most pressing environmental challenges of the 21st century, which is



ensuring access to sufficient water supply for human beings and ecosystems," said Laurence C. Smith, co-author of the study and chair of UCLA's geography department. "I'm very excited about this discovery."

A noted authority on climate change and its affect on the Arctic, Smith has been trying for the past 20 years to figure out a way to estimate <u>river</u> <u>discharge</u> on Earth from <u>outer space</u>. The closest he had been able to get to the goal was an approach that employed satellite imagery but still required on-site verification.

"You still needed measurements on the ground to calibrate the method, which defeated the purpose," he said. "But now we can estimate discharge with river width alone, and width we can see from space."

For the *PNAS* proof-of-concept study, Gleason and Smith used their new system and satellite imagery from the U.S. government's Landsat program to measure the discharge of three rivers—the Mississippi, Canada's Athabasca and China's Yangtze. They then compared those results with discharge figures from respected government agencies that monitor those rivers and whose calculations are recognized as highly accurate. The estimates of the UCLA geographers came within 20 to 30 percent of those figures.

"That might sound like a big error, but right now, we have no idea about the flow rates of most rivers around the world," Smith said. "To get a number that is within even 30 percent of accurate is incredibly helpful."

Smith tapped Gleason for field work on the <u>ice sheet</u> because Gleason had a track record of developing innovative remote-sensing applications. In 2012, for instance, he created an algorithm for calculating the number of tree crowns in a forest using aerial laser-scanning technology.

Greenland's ice sheet, the world's second largest body of ice after the



Antarctic ice sheet, originally covered 660,000 square miles, but it is melting at a rapid rate due to climate change. Runoff from the sheet is expected to elevate sea levels dramatically, and Smith is using field measurements in an attempt verify current computer-model-based projections of that rise.

Tasked with calculating the discharge rates of rivers carrying melt from the ice sheet, Gleason struggled during his first two summers in the field. With banks that act like quicksand and rapidly moving water carrying ice chunks the size of cars, the ice sheet's more remote rivers proved too dangerous to approach, much less enter and take <u>measurements</u>.

"We couldn't even get within 10 feet of a river—and at that, it was pushing it," Gleason said.

Moreover the rivers crisscross each other in a braided fashion that complicates calculation. Even instruments for measuring depth were useless; the rivers were too full of sediment and so turbulent they tossed them right out of the water.

However, Gleason was able to methodically photograph rivers from both sides and later use the information to calculate the waterways' widths. Back at UCLA, he pored over literature on the formula that had been used for the past 60 years to calculate river discharge. Gleason estimates that he read more than 100 studies on hydrologic geometry. Against all odds, his efforts uncovered a redundancy in the time-honored formula.

"Larry didn't believe me at the beginning, so he kept saying, 'OK, test it with these data and those data,' " Gleason recalls with a laugh.

Gleason's discovery allowed him to streamline the equation so that it required just one input —width—instead of three. The approach tracks vacillations in width at precise intervals on a river over time. Smith still



shakes his head in disbelief at the discovery.

"It's pretty amazing for this dusty old field that nobody discovered it before," Smith said.

Since submitting the article for publication, Gleason and Smith have tested the method on an additional 19 rivers worldwide. So far, the technique has attained at least the same level of accuracy as reported in *PNAS*, with the exception of one particular topology: The geographers found that the approach doesn't work on rivers with vertical concrete sides, like the river that runs through Los Angeles.

Ultimately, the geographers plan to test the method on a total of more than 30 rivers around the globe. With at least four per continent, the test rivers are being selected for the diversity of conditions that they represent.

"We're halfway done and the approach is continuing to work, and I've made it run faster and more accurately than in the *PNAS* paper," Gleason said.

And what of the Greenland rivers? When Gleason returned last year, he happened to see a herd of caribou crossing the waterway that had given him so much trouble. Based on where the water level hit the animals, he could extrapolate the depth measurement that had eluded him the two prior summers. Together with Smith, he improvised an approach to measuring velocity by throwing wadded balls of willow into the river and timing the balls' progress downstream. Finally, Gleason had the information to be able to figure out the river's discharge using the conventional approach.

As excited as he was, the achievement rang somewhat hollow.



"It was great to finally get the measurement, but by that time I was so hooked on the discovery that resulted from the work-around," he said. "A lot can be done with this new algorithm that we developed."

**More information:** Toward global mapping of river discharge using satellite images and at-many-stations hydraulic geometry, *PNAS*, <u>www.pnas.org/cgi/doi/10.1073/pnas.1317606111</u>

Provided by University of California, Los Angeles

Citation: Geographers create 'easy button' to calculate river flows from space (2014, March 18) retrieved 24 May 2024 from <u>https://phys.org/news/2014-03-geographers-easy-button-river-space.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.