

# 'Tree of life' distances are no shortcut to conservation

March 11 2014

---

Evolutionary distances that conservationists use to identify and target distinct species may be unreliable, Oxford University research suggests.

Some [conservation strategies](#) assume that the evolutionary distances between [species](#) on a phylogenetic 'tree of life' (a branching diagram of species popularised by Charles Darwin) can be used to predict how diverse their biological features will be. These distances are then used to select which species to conserve in order to maximise interesting biological features – such as potentially useful drug compounds and resilience to climate change.

But a new analysis of data from 223 studies of animals, plants, and fungi, shows that methods based on such distances are often no better at conserving interesting biological features than picking species at random. A report of the research is published this week in the journal *Diversity and Distributions*.

'Whilst 'close neighbours' on the branches of the tree of life are likely to share more biological features than distant ones, we found that you only have to move a short distance away before predictions about how much more diverse an organism's features should be are no better than a random choice,' said Dr Robert Scotland of Oxford University's Department of Plant Sciences. 'Much of this may be down to parallel or convergent evolution that sees similar biological features – such as eyes and wings – evolving independently again and again throughout the history of life.'

The new analysis suggests that phylogenetic distance by itself is not an adequate way of prioritising which organisms are most dissimilar to target for conservation.

'Maximising biological feature diversity is clearly important to conservation but you won't achieve this if you don't select the right range of species, and our study shows that you are unlikely to select the right range of species if you use phylogenetic distance,' said Dr Scotland. 'What our work suggests is that we need better, more nuanced, methods for identifying feature diverse species to underpin conservation strategies.'

**More information:** Kelly, S., Grenyer, R., Scotland, R. W. (2014), Phylogenetic trees do not reliably predict feature diversity. *Diversity and Distributions*. [DOI: 10.1111/ddi.12188](https://doi.org/10.1111/ddi.12188)

Provided by Oxford University

Citation: 'Tree of life' distances are no shortcut to conservation (2014, March 11) retrieved 12 August 2024 from <https://phys.org/news/2014-03-tree-life-distances-shortcut.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.