

New wireless network to revolutionise soil testing

March 13 2014

A University of Southampton researcher has helped to develop a wireless network of sensors that is set to revolutionise soil-based salinity measuring.

Dr Nick Harris, from Electronics and Electrical Engineering, worked with a group of professors from the University of Western Australia (UWA) to produce the revolutionary sensor that can carry out non-destructive testing of [soil](#) samples.

The sensor is capable of measuring the [chloride](#) (salt) in the soil moisture and linking up with other [sensors](#) to create a wireless network that can collate and relay the measurement readings. The network can also control the time intervals at which measurements are taken.

The sensor is placed in the soil and measures the chloride levels in the soil moisture in a non-destructive way. These [chloride levels](#) make up a high proportion of the overall soil salinity.

Dr Harris says: "Traditionally, soil-based measurements involve taking samples and transporting them to the laboratory for analysis. This is very labour and cost intensive and therefore it usually means spot checks only with samples being taken every two to three months. It also doesn't differentiate between chloride in crystallised form and chloride in dissolved form. This can be an important difference as plants only 'see' chloride in the [soil moisture](#).

"The removal of a soil sample from its natural environment also means that the same sample can only be measured once, so the traditional (destructive) method is not suited to measuring changes at a point over a period of time.."

The new sensors are connected to a small unit and can be 'planted' in the ground and left to their own devices. The limiting factor for lifetime is usually the sensor. However, these sensors are expected to have a lifetime in excess of one year. The battery-powered unit can transmit data and information by short range radio, Bluetooth, satellite or [mobile phone network](#), as well as allowing data to be logged to a memory card to be collected later.

The novel device allows up to seven sensors to be connected at a time to a single transmitter allowing multi-point measurements to be simply taken.

Dr Harris adds: "These soil-based chloride sensors can benefit a wide range of applications. Large parts of the world have problems with salt causing agricultural land to be unusable, but the new sensors allow the level of salt to be measured in real time, rather than once every few months as was previously the case.

"At plant level, probes can be positioned at continuous levels of depth to determine the salt concentration to which roots are exposed and whether this concentration changes with the depth of the soil or in different weather conditions. We can also measure how well a plant performs at a particular concentration and change the salt content for a few days and observe the effects.

"On a bigger scale, sensors could be placed at different locations at catchment scale to observe any changes in the level of salinity within a field over time, having a direct impact on irrigation strategies. We have

already been able to make some interesting observations on real world chloride concentration changes over just 24 hour periods, illustrating the dangers of relying on single point, single time measurements."

Provided by University of Southampton

Citation: New wireless network to revolutionise soil testing (2014, March 13) retrieved 28 April 2024 from <https://phys.org/news/2014-03-wireless-network-revolutionise-soil.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--