

Arid areas absorb unexpected amounts of atmospheric carbon

April 6 2014



Mojave Desert plots like this see from a radio-controlled helicopter showed researchers that arid areas can absorb unexpected amounts of atmospheric carbon. Credit: Desert Research Institute/ University of Nevada at Las Vegas

Researchers led by a Washington State University biologist have found that arid areas, among the biggest ecosystems on the planet, take up an unexpectedly large amount of carbon as levels of carbon dioxide increase in the atmosphere. The findings give scientists a better handle on the earth's carbon budget—how much carbon remains in the atmosphere as CO₂, contributing to global warming, and how much gets stored in the land or ocean in other carbon-containing forms.

"It has pointed out the importance of these arid ecosystems," said R. Dave Evans, a WSU professor of biological sciences specializing in ecology and global change. "They are a major sink for atmospheric [carbon](#) dioxide, so as CO₂ levels go up, they'll increase their uptake of CO₂ from the atmosphere. They'll help take up some of that excess CO₂ going into the atmosphere. They can't take it all up, but they'll help."

The findings, published in the journal *Nature Climate Change*, come after a novel 10-year experiment in which researchers exposed plots in the Mojave Desert to elevated [carbon-dioxide](#) levels similar to those expected in 2050. The researchers then removed soil and plants down to a meter deep and measured how much carbon was absorbed.

"We just dug up the whole site and measured everything," said Evans.

The idea for the experiment originated with scientists at Nevada's universities in Reno and Las Vegas and the Desert Research Institute. Evans was brought in for his expertise in nutrient cycling and deserts, while researchers at the University of Idaho, Northern Arizona University, Arizona State University and Colorado State University also contributed.

Funding came from the U.S. Department of Energy's Terrestrial Carbon Processes Program and the National Science Foundation's Ecosystem Studies Program. The work addresses one of the big unknowns of [global](#)

warming: the degree to which land-based ecosystems absorb or release carbon dioxide as it increases in the atmosphere.



Washington State University researchers Ben Harlow, left, and Dave Evans found soil from arid areas take up an unexpectedly large amount of carbon as levels of carbon dioxide increase in the atmosphere. Credit: Shelly Hanks, Washington State University


Receiving less than 10 inches of rain a year, arid areas run in a wide band at 30 degrees north and south latitude. Along with semi-arid areas, which receive less than 20 inches of rain a year, they account for nearly half the earth's land surface.

Forest soils have more organic matter and, square foot for square foot, hold much more carbon. But because arid soils cover so much area, they

can have an outsized role in the earth's carbon budget and in how much the earth warms as heat-trapping gases accumulate in the atmosphere.

Working on the Nevada National Security Site, the researchers marked off nine octagonal plots about 75 feet in diameter. Air with 380 parts-per-million concentrations of CO₂, the current CO₂ levels, was blown over three plots. Three received no extra air. Three were exposed to concentrations of 550 parts per million, the CO₂ levels expected in 2050.

The CO₂ was fed through PVC pipes ringing the plots and had a specific chemical fingerprint that could be detected when the soil, plants and other biomass were analyzed.



Akihiro Koyama, now of Colorado State University, and Ben Harlow of Washington State University gather soil samples from a test plot in the Mojave Desert. Credit: Desert Research Institute

The analysis, done by Benjamin Harlow in WSU's Stable Isotope Core Laboratory, suggests that arid lands may increase their carbon uptake enough in the future to account for 15 to 28 percent of the amount currently being absorbed by land surfaces.

Overall, said Evans, rising CO₂ levels may increase the uptake by [arid lands](#) enough to account for 4 to 8 percent of current emissions.

The experiment did not account for other possible changes stemming from climate change, like varying precipitation and warming temperatures.

Still, said Evans, "I was surprised at the magnitude of the carbon gain, that we were able to detect it after 10 years, because 10 years isn't very long in the life of an ecosystem."

While forest ecosystems tend to store carbon in plant matter, the Mojave researchers found most carbon was being taken up by increased activity in the rhizosphere, a microorganism-rich area around the roots.

From an optimistic point of view, the research suggests that, come 2050, [arid ecosystems](#) will be doing more than their fair share of taking earth-warming carbon out of the atmosphere. But a potential cause for concern is what happens to these ecosystems as the planet's population grows and people look for places to develop and live.

"Land is extremely valuable," said Evans. "A lot of growth may occur in these areas that are fairly arid and we don't know what that's going to do then to the [carbon budget](#) of these systems."

More information: Paper: Greater ecosystem carbon in the Mojave

Desert after ten years exposure to elevated CO₂, [DOI: 10.1038/nclimate2184](#)

Provided by Washington State University

Citation: Arid areas absorb unexpected amounts of atmospheric carbon (2014, April 6) retrieved 16 August 2024 from <https://phys.org/news/2014-04-arid-areas-absorb-unexpected-amounts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.