

Atmospheric boundary layer exacerbated mega heat waves

April 21 2014, by Dr.ir. Aj (Ryan) Teuling



Vegetation on dry soils, such as in this picture from Australia, have limited water available for evaporation. Because of reduced evaporation, heat can accumulate in the lowest part of the atmosphere, the so-called boundary layer. Through this process, extreme heatwaves can occur during drought conditions in regions that are normally humid.

The extreme nature of the heat waves of 2003 in Western Europe and of

2010 in Russia and Eastern Europe even surprised scientists at the time. NWO Veni researcher Ryan Teuling from Wageningen University says that the extreme temperatures can be explained by the interaction between dry soils with the atmospheric boundary layer - the lowest part of the atmosphere. The role of this boundary layer has received too little attention in studies using existing weather models, he claims in *Nature Geoscience*.

The mega [heat wave](#) of August 2003 in Western Europe broke various temperature records at the time, with temperatures of 40°C in France. The economic damage was estimated at between five and ten billion euros, due to forest fires, air pollution and loss of agricultural productivity. In Paris alone, thousands died as a consequence of the high temperatures. Researchers were unfamiliar with such heat waves in Europe and thought that it was a one-off, exceptional event. At least that was the case until 2010 when new records were set, this time in Eastern Europe and Russia.

Atmospheric boundary layer

In case of an atmospheric blocking situation, soil desiccation and a build-up of heat in the atmospheric [boundary layer](#) all occur at the same time, then exceptional heat waves are possible. That is the conclusion of Dr Ryan Teuling, assistant professor in hydrology and quantitative water management in Wageningen. The unusual atmospheric situation had previously been demonstrated to be the most important cause of the two heat waves. A high-pressure area blocked the penetration of low-pressure areas with colder air and wind. The researchers demonstrated that for both heat waves the same developments in the boundary layer conditions occurred in combination with an enhanced interaction between soil desiccation and heat accumulation in the atmospheric boundary layer. This lowest part of the troposphere is several tens to hundreds of metres thick at night, and up to several kilometres thick

during the day.



Over a period of several days to several weeks heat was built up in an increasingly thicker atmospheric boundary layer, helped by heat convection from more southerly situated warm areas. During this period the soil dried out even further, leading to a reduced cooling effect of water evaporation from the surface. This process was enhanced by the [high temperatures](#). Due to cooling at night, the warm air layer became decoupled from the earth's surface, as a result of which the heat of the following day could lead to increasingly higher temperatures. The blanket of warm air above the earth's surface became increasingly thicker during the heat wave.

45 degrees Celsius

Teuling: 'With this new knowledge we can make predictions about possible temperatures during heat waves. Temperatures that gradually build up to 40°C are indeed possible in this part of the world. However, sudden [temperature](#) rises to 45°C as occurred during heat waves in, for example, Melbourne are unlikely here. Therefore the distance to the Equator and to large desert areas is too great.'

The researchers made use of satellite data and measurements from weather balloons of the situation in France in 2003 and Russia in 2010. They combined these data with a coupled soil-water-atmosphere model.

Provided by Wageningen University

Citation: Atmospheric boundary layer exacerbated mega heat waves (2014, April 21) retrieved 20 April 2024 from

<https://phys.org/news/2014-04-atmospheric-boundary-layer-exacerbated-mega.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.