

Landmark experiment on wave interference from the early 1800s is revisited using gold nanoparticles

April 9 2014



Example of the energy flow and optical vortices found around closely spaced gold nanoparticles. The effects resemble the field lines seen in Young's slit experiments. Credit: A*STAR Data Storage Institute

In the eighteenth century, scientists faced a conundrum: is light a wave or a particle? One of strongest pieces of evidence to support the 'wave view'—the landmark double-slit experiment—was reported in 1804 by the scientist Thomas Young. Young passed coherent light through two closely spaced slits and observed a set of interference fringes, a result



that occurs with wave phenomena like sound or water. This observation became the basis for the modern wave theory of light.

Two hundred years later, Arseniy Kuznetsov and co-workers from the A*STAR Data Storage Institute, together with collaborators in Australia, Singapore, the United Kingdom and Russia, have performed an experiment analogous to Young's experiments but using nanoscale objects. The team studied the light scattering in the visible and near-infrared wavelength regions from a cluster of two or three closely spaced gold plasmonic nanoparticles. They observed interference and resonance effects that resemble those seen in Young's experiments.

In particular, while studying a trimer system consisting of three discrete metallic nanodisks of about 145 nanometers in diameter and 60 nanometers thick, the team found evidence for the presence of near-field, subwavelength-sized optical vortices and the circulation of electromagnetic energy (see image). This finding is very similar to what occurs to the <u>energy flow</u> pattern in a Young-type experiment performed with three slits.

One of the key issues in nanoplasmonics is the interaction between metallic nanoparticles at the nanoscale. "Even if the separation between two or multiple non-periodically arranged nanoparticles is of the order of wavelength, their interaction can be strong enough to change their scattering and absorption properties," notes Kuznetsov. "This can be explained by the peculiarities of the Poynting vector (energy) flow around the nanoparticles and formation of optical vortices, which produce a pattern of field lines similar to Young's classic experiment."

The team's findings, says Kuznetsov, not only expand our fundamental understanding of how light interacts with nanoclusters of metallic particles, but have both theoretical and practical applications. "They may also prove useful for applications such as improved solar cells and



plasmonic biosensors." However, their most remarkable application, he suggests, may be in the emerging area of nanoantennas.

In the future, the team is aiming to study the resonant properties and interactions of nanoparticles made from nonmetallic materials. In particular, they plan to investigate high-refractive index dielectric materials such as silicon, which, unlike <u>metallic particles</u>, do not suffer from high optical losses.

More information: Rahmani, M., Miroshnichenko, A. E., Lei, D. Y., Luk'yanchuk, B., Tribelsky, M. I. et al. "Beyond the hybridization effects in plasmonic nanoclusters: Diffraction-induced enhanced absorption and scattering." *Small* 10, 576–583 (2013). <u>dx.doi.org/10.1002/smll.201301419</u>

Provided by Agency for Science, Technology and Research (A*STAR), Singapore

Citation: Landmark experiment on wave interference from the early 1800s is revisited using gold nanoparticles (2014, April 9) retrieved 1 May 2024 from <u>https://phys.org/news/2014-04-landmark-early-1800s-revisited-gold.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.