


# Organic solar cells more efficient with molecules face-to-face

April 7 2014, by Tracey Peake

---



This shows molecules in face-on orientation inside organic solar cell. Credit: NC State University/ Peter Allen

(Phys.org) —New research from North Carolina State University and UNC-Chapel Hill reveals that energy is transferred more efficiently

inside of complex, three-dimensional organic solar cells when the donor molecules align face-on, rather than edge-on, relative to the acceptor. This finding may aid in the design and manufacture of more efficient and economically viable organic solar cell technology.

Organic [solar cell efficiency](#) depends upon the ease with which an exciton – the energy particle created when light is absorbed by the material – can find the interface between the donor and [acceptor](#) molecules within the cell. At the interface, the exciton is converted into charges that travel to the electrodes, creating power. While this sounds straightforward enough, the reality is that molecules within the donor and acceptor layers can mix, cluster into domains, or both, leading to variances in domain purity and size which can affect the power conversion process. Moreover, the donor and acceptor molecules have different shapes, and the way they are oriented relative to one another matters. This complexity makes it very difficult to measure the important characteristics of their structure.

NC State physicist Harald Ade, UNC-Chapel Hill chemist Wei You and collaborators from both institutions studied the molecular composition of [solar cells](#) in order to determine what aspects of the structures have the most impact on efficiency. In this project the team used advanced soft X-ray techniques to describe the orientation of molecules within the donor and acceptor materials. By manipulating this orientation in different solar cell polymers, they were able to show that a face-on alignment between donor and acceptor was much more efficient in generating power than an edge-on alignment.

"A face-on orientation is thought to allow favorable interactions for charge transfer and inhibit recombination, or charge loss, in [organic solar cells](#)," Ade says, "though precisely what happens on the molecular level is still unclear.

"Donor and acceptor layers don't just lie flat against each other," Ade explains. "There's a lot of mixing going on at the molecular level. Picture a bowl of flat pasta, like fettuccine, as the [donor](#) polymer, and then add 'ground meat,' or a round acceptor molecule, and stir it all together. That's your solar cell. What we want to measure, and what matters in terms of efficiency, is whether the flat part of the fettuccine hugs the round pieces of meat – a face-on orientation – or if the fettuccine is more randomly oriented, or worst case, only the narrow edges of stacked up pasta touch the meat in an edge-on orientation. It's a complicated problem.

"This research gives us a method for measuring this molecular orientation, and will allow us to find out what the effects of orientation are and how [orientation](#) can be fine-tuned or controlled."

The paper appears online April 6 in *Nature Photonics*.

**More information:** "The influence of molecular orientation on organic bulk heterojunction solar cells" John R. Tumbleston, Brian A. Collins, Eliot Gann, Wei Ma and Harald Ade, North Carolina State University; Liqiang Yang, Andrew C. Stuart and Wei You, University of North Carolina at Chapel Hill, Published: April 6, 2014, in *Nature Photonics*

### **Abstract**

In bulk heterojunction organic photovoltaics, electron-donating and electron-accepting materials form a distributed network of heterointerfaces in the photoactive layer, where critical photo-physical processes occur. However, little is known about the structural properties of these interfaces due to their complex three-dimensional arrangement and the lack of techniques to measure local order. Here, we report that molecular orientation relative to donor/acceptor heterojunctions is an important parameter in realizing high-performance fullerene-based, bulk

heterojunction solar cells. Using resonant soft X-ray scattering, we characterize the degree of molecular orientation, an order parameter that describes face-on (+1) or edge-on (-1) orientations relative to these heterointerfaces. By manipulating the degree of molecular orientation through the choice of molecular chemistry and the characteristics of the processing solvent, we are able to show the importance of this structural parameter on the performance of bulk heterojunction organic photovoltaic devices featuring the electron-donating polymers PNDT–DTBT, PBnDT–DTBT or PBnDT–TAZ.

Provided by North Carolina State University

Citation: Organic solar cells more efficient with molecules face-to-face (2014, April 7) retrieved 5 May 2024 from

<https://phys.org/news/2014-04-solar-cells-efficient-molecules-face-to-face.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.