


Researchers track ripples in freestanding graphene for first time

April 30 2014, by Chris Branam



Paul Thibado, University of Arkansas professor of physics. Credit: Russell Cothren

(Phys.org) —An international team of scientists, led by physicists at the University of Arkansas, has tracked the dynamic movement of ripples in freestanding graphene at the atomic level.

This discovery advances the fundamental understanding of one of the

strongest, lightest and most conductive materials, said Paul Thibado, University of Arkansas professor of physics.

"Physicists have known that the [ripples](#) must be there and some experiments did find them," he said. "But they could only measure the ripples as static in time. The theory requires that they fluctuate, more like looking at an ocean with waves. The thermal energy needs to vibrate. Up until our experiment no one had successfully measured this dynamic property of the ripples."

The team published its findings on Monday, April 28, in *Nature Communications*, an online journal published by the journal *Nature*, in a paper titled "Unusual ultra-low frequency fluctuations in freestanding graphene."

Freestanding graphene could emerge as a replacement for silicon and other materials in microprocessors and next-generation energy devices, but much remains unknown about its mechanical and thermal properties.

Graphene, discovered in 2004, is a one-atom-thick sheet of graphite. Electrons moving through graphite have mass and encounter resistance, but electrons moving through graphene are massless and therefore encounter much less resistance. This makes graphene an excellent candidate material for future energy needs, as well as for use in quantum computers, to enable enormous calculations with little energy use.

The study was led by Peng Xu, a postdoctoral research associate in the department of physics in the J. William Fulbright College of Arts and Sciences at the University of Arkansas.

Xu and Thibado used scanning tunneling microscopy, which produces images of individual atoms on a surface, to measure ultra-low frequency fluctuations in a one-square-angstrom region of freestanding graphene.

An angstrom is a unit of length equivalent to one hundred millionth of a centimeter.

These fluctuations, known as intrinsic ripples, have been exceedingly difficult to study because their vertical movement usually creates blurry images, Thibado said. The University of Arkansas researchers successfully produced clear images, enabling them to present a model from elasticity theory to explain the very-low frequency oscillations. In physics, elasticity is the tendency of solid materials to return to their original shape after being deformed.

The researchers' innovative scanning tunneling microscopy technique provides a much-needed atomic- scale probe for the time-dependent behaviors of intrinsic ripples, said Thibado, an expert in experimental [condensed matter physics](#). The ripple dynamics are important for understanding mechanical stability and the efficient thermal conductivity transport properties of graphene.

In the last decade, theoretical physicists predicted a bending mode in two-dimensional material graphene that couples to a stretching mode of the graphene. Without that bending and coupling, freestanding [graphene](#) wouldn't exist, Thibado said.

More information: "Unusual ultra-low-frequency fluctuations in freestanding graphene." P. Xu, et al. *Nature Communications* 5, Article number: 3720 [DOI: 10.1038/ncomms4720](https://doi.org/10.1038/ncomms4720). Received 09 September 2013 Accepted 26 March 2014 Published 28 April 2014

Provided by University of Arkansas

Citation: Researchers track ripples in freestanding graphene for first time (2014, April 30)

retrieved 19 April 2024 from

<https://phys.org/news/2014-04-track-ripples-freestanding-graphene.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.