


Far out: A giant exoplanet where none has been seen before

May 23 2014, by Sheyna E. Gifford, Md



Timeline of where planets are expected to be found as their parent stars evolve when forming in a protoplanetary disk. The star GU Psc is part of the AB Doradus Moving Group, with an estimated age between 70-130 million years. As such, GU Psc b would be expected to be within 100 AU of GU Psc. Instead, it is 2000 AU away, suggesting a different formation mechanism. Credit: Shu et al. 1987, Courtesy of the James Webb Telescope and NASA

Humans have an eye for the familiar: for people, for civilizations, for planets and planetary systems that match what we have seen in the past. For this reason, as well as a few others, we rarely find something truly



unique in the universe. When we do, it's often by happenstance.

Finding a planet ten times the <u>mass</u> of Jupiter orbiting 40 times further out than Pluto was in some ways unexpected.


"It's not a type of exoplanet we had previously found and it's certainly not one that the theorists were expecting," said paper co-author PhD student Marie-Eve Naud at the University of Montréal, "especially not around low-mass stars like GU Psc (~30% the mass of the Sun)."

Last June, Astrobio.net ran the surprising story of a planet 5-10 times the mass of Jupiter that formed 80 AU from its <u>parent star</u>. In yet another twist in the continuing saga of planetary treasure-hunting, researchers in Naud's group at the University of Montréal announced the discovery of a planet many times the mass of Jupiter at an incredible distance from its star: 2000 AU.

This planet-GU Psc b-is so far from its star that it could be photographed without the aid of adaptive optics, appearing as an independent point of infrared light from its star GU Psc.

Unlike GU Psc b, most of the 1700 plus exoplanets in NASA's current catalog have not been directly imaged, or photographed directly, in any wavelength. Their gravitational effects on their parent stars have been measured. Sometimes the dimming effect as they occlude the light from their parent stars have been recorded. Occasionally, existence of planet is inferred by the presence of a gap in dust disks rotating around a star.





Credit: © Lucas Granito

At present, the Exoplanet catalog maintained by the Paris Observatory lists only 46 <u>planets</u> that have been directly imaged, of which GU Psc b is unique in a number of ways.

Of the exoplanets that have been photographed, GU Psc b at 9-13 times the mass of Jupiter is one the largest that is probably not a failed star, or brown dwarf. At thirteen times the mass of Jupiter, a gas giant attains the minimum mass to begin some nuclear fusion early in its life. Many of the photographed companions might in fact have a mass greater than 13 times that of Jupiter, and may be more like brown dwarfs than planets in nature. GU Psc b is also one of the few so far out from its companion. Perhaps most interestingly, of all the far-out planets discovered so far by any means, this new planet is only giant, distant one at all that rotates around a star like our own.




"GU Psc b is certainly the farthest planetary-mass companion known around a regular, main-sequence star," said Naud.

In their recent paper, Naud and her co-advisor at the University of Montreal Dr. Étienne Artigau describe how they applied an old technique in a new way. Seeing giant planets that are relatively close to their stars, many light years away, requires state-of-the art adaptive optics (AO) and the Gemini Observatory. Naud and Artigau had the idea of looking for farther out planets using a "standard" camera. To find distant planets without specialized instruments, they exploited certain color, or wavelengths, that distinguish planets from stars or galaxies in the background. They surveyed dozens of stars with this technique before finding GU Psc b.

"My idea with this project was to explore the entire 'sphere' where a planet could physically stay for an astrophysically significant amount of time," said Artigau, "regardless of predictions relative to formation mechanisms."

Many models predict that planets should not be found beyond 100 AU. Those same models also predict that if planets are present within about 5000 AU of a parent star, they should stay there for billions of years. For astronomers, that means there is a wide swath of sky that can potentially contain a planet. Regardless, many recent searches have focused on finding planets very near young stars. Naud and her colleagues took a slightly broader view.





Credit: ©Stellarium

"This does not allow us to find planets as close as the ones found by specialized

instruments, but it allowed us to detect GU Psc b, located 42 arc seconds from its star," said Naud, "GU Psc b is precious to validate our models of similar objects, and we will be able to use the information we obtain on this object to better understand other, closer-in and harder to study planetary-mass companions.

Because GU Psc b is so far from its parent star, astronomers can learn a tremendous amount about it. In just one hour on the Gemini North scope, Naud and her team obtained a spectrum of GU Psc b that allowed them to estimate important physical properties of the planet, like mass and temperature. These are usually difficult or impossible to observe because of the nearby star.



As a very distant companion, GU Psc b is a diamond in the rough. Not only does it provide us a very clear view of the planet itself with its mass and temperature, but it also provides us with an impetus to look for more planets like it, in places we would have never looked before.

"This type of object exists, and we can quite easily find them," said Naud, "so we should definitely search for them."

More information: Study paper: <u>iopscience.iop.org/0004-637X/7</u>... <u>004-637X_787_1_5.pdf</u>

Provided by Astrobio.net

Citation: Far out: A giant exoplanet where none has been seen before (2014, May 23) retrieved 27 April 2024 from <u>https://phys.org/news/2014-05-giant-exoplanet.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.