

# Scientists reveals how fish were able to colonise poisonous springs

May 12 2014

---



*Poecilia mexicana*. Credit: M.Pfenninger

Hydrogen sulphide ( $H_2S$ ) is a potent inhibitor of aerobic respiration. However populations of shortfin molly fish managed to colonise springs with high concentrations of dissolved hydrogen sulphide. In a new study researchers from LOEWE Biodiversity and Climate Research Centre (BiK-F) and the Goethe University Frankfurt am Main present evidence

of genetic changes minimizing the harmful effects of H<sub>2</sub>S which enable the fish to survive in this deleterious environment. The study provides insight into the molecular mechanisms of this key adaptation for the first time. It is published online today in *Nature Communications*.

Shortfin molly fishes (*Poecilia mexicana*) may only measure a few inches, but they are still exceptional. Populations of *Poecilia mexicana*, whose relatives are the well-known guppy, colonised sulphide-rich volcanic springs in Southern Mexico. In making this particular habitat their home, they have made the impossible possible, because [hydrogen sulphide](#) (H<sub>2</sub>S), as for many other animal, is lethal. Even at low concentrations the gas blocks the cytochrome c oxidase-complex (COX). The higher the level of hydrogen sulphide, the more the activity of COX is inhibited. As it is essential for respiration, this turns out to be lethal in the end.

## **Changes in genetic make-up make less susceptible to poison**

A team led by Prof. Dr. Markus Pfenninger, LOEWE Biodiversity and Climate Research Centre (BiK-F) and PD Dr. Martin Plath, Goethe University, has taken a closer look at the survivors. Their analysis showed that the COX activity of individuals of shortfin molly fish which colonise H<sub>2</sub>S-rich waters remains virtually unchanged under high H<sub>2</sub>S concentrations. This is due to number of changes in the *cox1* and *cox3* genes, which have only occurred in populations living in the poisonous springs. Thus, transplanting individuals from non-sulphidic habitat to springs with high H<sub>2</sub>S levels kills them for sure.

## **Molecular mechanisms of adaptation to extreme habitat**

"In this paper we analyse the key adaptation to an extreme habitat up to its molecular basis at the level of amino acids. This way, for the first time, we are able to point out, where exactly the adaptation has taken place." Pfenninger concludes. The team also modelled three dimensional protein structures in order to shed light on necessary significant structural changes of amino acids in the *cox1* gene. Without these structural changes, the colonisation of the H<sub>2</sub>S-containing water for the fish would have been impossible. By colonising the poisonous springs, where there are hardly any other competitors, the fish may feed on resistant midge larvae that also occur there.

## **Closely related fish follow different paths to adaptation**

The study also shows that closely related populations of a species follow parallel as well as disparate paths in response to similar environmental conditions. Three shortfin molly fish populations were sampled for study. Two of the populations show the same changes in their genetic material in adapting to the hostile conditions. However this proved to be not the case for the third population of shortfin molly fish. Whereas these fish also tolerate high levels hydrogen sulphide, the mechanism enabling their adaptation is still subject to ongoing research.

**More information:** Pfenninger, M. et al.: "Parallel evolution of *cox*-genes in H<sub>2</sub>S- tolerant fish as key adaptation to a toxic environment" *Nature Communications*, [DOI: 10.1038/ncomms4873](https://doi.org/10.1038/ncomms4873)

Provided by Senckenberg Research Institute and Natural History Museum

Citation: Scientists reveals how fish were able to colonise poisonous springs (2014, May 12)  
retrieved 22 April 2024 from

<https://phys.org/news/2014-05-scientists-reveals-fish-colonise-poisonous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.