

Litter-dwelling thrips live mainly in tropical and subtropical regions

June 18 2014



This is an example of *Psalidothrips ascitus (Ananthakrishnan)*. Abundant individuals were found in leaf litter. Credit: Dr Jun Wang

The species diversity in soil fauna has been studied in temperate regions for more than 50 years, but with scarcely any mention of thrips. This lack of reference to thrips raises the question whether or not litter-



dwelling thrips are distributed only in tropical and sub-tropical regions.

To answer this question a total 150 <u>leaf litter</u> samples were collected from 6 natural reserves located in three climatic regions, temperate, subtropical and tropical, along a 4100 km latitudinal gradient in East China. The survey was done over a four-year period by Dr Jun Wang, who is a thrips specialist and an assistant professor at the College of Plant Science, Jilin University, China. His results have been published in the open access journal *ZooKeys*

'Thrips constitute over 3.0% of total litter–dwelling macroinvertebrate individuals in 4 natural reserves from subtropical and tropical zone', said Dr Wang. 'In contrast, it constitute only 0.3% in the warm temperate natural reserves, and no thrips is collected in a mid temperate reserve.'

Dr Wang said that the order on the average species numbers per plot of litter thrips was tropic followed by subtropics followed by temperate. Mean density of litter thrips per plots in the tropics and subtropics was significantly higher than that in the temperate region.

Dr. Laurence Mound, CSIRO Entomology, Australia commented that this manuscript is the first serious attempt to look at the diversity of thrips in leaf-litter, based on a good sampling strategy, and investigating the variation with latitude.

More information: Wang J, Tong X, Wu D (2014) The effect of latitudinal gradient on the species diversity of Chinese litter-dwelling thrips. *ZooKeys* 417: 9. DOI: 10.3897/zookeys.417.7895

Provided by Pensoft Publishers



Citation: Litter-dwelling thrips live mainly in tropical and subtropical regions (2014, June 18) retrieved 20 April 2024 from

https://phys.org/news/2014-06-litter-dwelling-thrips-tropical-subtropical-regions.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.