

Sharpening a test for tracing food-borne illness to source

June 23 2014

Research from the University of Melbourne, Australia, could make it easier for public health investigators to determine if a case of food poisoning is an isolated incident or part of a larger outbreak. The findings are published ahead of print in the *Journal of Bacteriology*.

The study focuses on a test called multi-locus variable number tandem repeats variable analysis (MLVA). The test, which is increasingly used in the detection and investigation of foodborne outbreaks, analyzes specific sequences of DNA (called loci) that change rapidly enough over time to distinguish [outbreak](#) strains from other circulating strains of the bacteria but not so rapidly that connections could be masked by changes arising during the course of an outbreak.

However, the rates at which MLVA profiles change have not been directly investigated for *Salmonella*, and thus it is sometimes unclear how these profiles should be interpreted in the context of outbreak detection and investigation.

In the study, the investigators grew an isolate of *Salmonella Typhimurium* from an Australian [food poisoning](#) outbreak, and observed changes in its MLVA profile during more than 28,000 generations of growth in the laboratory. Then, using the same bacterial lineage, they observed changes in MLVA profile during 500 days of growth in mice.

They estimated the rates of copy number change at each of the five loci that are commonly used for *S. Typhimurium* MLVA. Three of the loci

saw changes in the DNA, but two did not. Based on these results, the researchers are recommending that isolates with zero or one variation in the three rapidly changing loci but no differences in the other two should be considered part of the same cluster.

They also noted that the relative rates of change among the loci were the same in the Petri dish studies and in the mouse study.

"This tells us we don't need to worry about where the bacteria were isolated from—humans or food," says Kathryn Holt, an author on the study.

MLVA is used for investigations of food-borne illnesses besides *Salmonella*, including *Listeria*, and *E. coli*. It is the primary method for investigations of *Salmonella* outbreaks in Europe, the UK, Australia, and elsewhere, says Holt.

"In the U.S., the Centers for Disease Control and Prevention uses another technique called PFGE for initial investigations and follows that with MLVA," she says.

Provided by American Society for Microbiology

Citation: Sharpening a test for tracing food-borne illness to source (2014, June 23) retrieved 25 June 2024 from <https://phys.org/news/2014-06-sharpening-food-borne-illness-source.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.