

South-west plants share fungal symbiosis for nitrogen uptake

June 11 2014, by Aaron Bryans



The Morrison Featherflower (*Verticordia nitens* (Lindl.) Endl.) was one of the south-west species used in the study. Credit: ron_n_beth

An investigation into nitrogen transfers between plants has found that different species can share nutrients through fungal interactions.

The study looked into numerous nutrient-acquisition strategies of plants to determine if nitrogen transfers between [native plants](#) exist in nutrient-poor soils in south-west Australia.

We've had this question [of] how is it possible that our ecosystem has so many species when the soils are so poor?" co-author Professor Erik Veneklaas says.

"What we've shown here is that it is the balance between positive and [negative interactions](#) that could help species survive together and in a way collaborate whilst they also compete."

Lead author and UWA's François Teste and his team found that plant species connected by both arbuscular mycorrhizal (AM) and ectomycorrhizal (EM) fungal interactions benefitted most from nitrogen transfer between plants.

AM interactions penetrate plant roots allowing nutrient intake, whilst EM interactions form sheaths around the roots of plants rather than infiltrating them.

"My work highlights that these interactions might also be partly responsible for biodiversity; so nutrient retention and sharing can be part of the reason why so many species are interacting together," Dr Teste says.

Nitrogen is an essential nutrient for plants as it is a basic component of chlorophyll that is used to process sunlight into energy through photosynthesis.

It also increases the strength and support of roots, allowing plants to take in more water and nutrients.

The transfer of nitrogen in various chemical forms between the atmosphere, the earth, plants and mammals is the [nitrogen cycle](#) and is essential for all forms of life.

Findings lend support for [restoration efforts](#)

The findings highlight the importance of plant diversity in ecosystems with Dr Teste hoping for improvements to restoration efforts.

"What I would wish is that in terms of restoration hopefully my work can highlight that assemblages and interactions are also part of the equation to successfully restoring areas that are having a hard time," Dr Teste says.

The exploration of fungal [interactions](#) between plants adds to prior knowledge of [plants](#) receiving and donating nitrogen through root transfers in addition to nitrogen obtained from the atmosphere as a result of post-lightning rain and [nitrogen](#) fixing bacteria.

Dr Teste and Professor Veneklaas are two of four WA scientists who conducted the study, the others being Professor Kingsley Dixon and Professor Hans Lambers.

More information: "Is nitrogen transfer among plants enhanced by contrasting nutrient-acquisition strategies?" Teste FP, Veneklaas EJ, Dixon KW, Lambers H. *Plant Cell Environ.* 2014 May 8. [DOI: 10.1111/pce.12367](#). [Epub ahead of print]

Provided by Science Network WA

Citation: South-west plants share fungal symbiosis for nitrogen uptake (2014, June 11) retrieved 25 April 2024 from <https://phys.org/news/2014-06-south-west-fungal-symbiosis-nitrogen-uptake.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.