


# Strange physics turns off laser

June 17 2014

---



Manipulating minute areas of gain and loss within individual lasers (shown as peaks and valleys in the image), researchers were able to create paradoxical interactions between two nearby lasers. Credit: Vienna University of Technology

Inspired by anomalies that arise in certain mathematical equations, researchers have demonstrated a laser system that paradoxically turns off when more power is added rather than becoming continuously brighter.


The finding by a team of researchers at Vienna University of

Technology and Princeton University, could lead to new ways to manipulate the interaction of electronics and light, an important tool in modern communications networks and high-speed information processing.

The researchers published their results June 13 in the journal *Nature Communications*<sup>1</sup>.

Their system involves two tiny lasers, each one-tenth of a millimeter in diameter, or about the width of a human hair. The two are nearly touching, separated by a distance 50 times smaller than the lasers themselves. One is pumped with electric current until it starts to emit light, as is normal for lasers. Power is then added slowly to the other, but instead of it also turning on and emitting even more light, the whole system shuts off.

"This is not the normal interference that we know," said Hakan Türeci, assistant professor of electrical engineering at Princeton, referring to the common phenomenon of light waves or sound waves from two sources cancelling each other. Instead, he said, the cancellation arises from the careful distribution of energy loss within an overall system that is being amplified.



An electron microscope image shows two lasers placed just two microns apart from each other. Credit: Vienna University of Technology

"Loss is something you normally are trying to avoid," Türeci said. "In this case, we take advantage of it and it gives us a different dimension we can use – a new tool – in controlling optical systems."

The research grows out of Türeci's longstanding work on mathematical models that describe the behavior of lasers. In 2008<sup>2</sup>, he established a mathematical framework for understanding the unique properties and complex interactions that are possible in extremely small lasers – devices with features measured in micrometers or nanometers. Different from conventional desk-top lasers, these devices fit on a computer chip.

That work opened the door to manipulating gain or loss (the

amplification or loss of an energy input) within a [laser system](#). In particular, it allowed researchers to judiciously control the spatial distribution of gain and loss within a single system, with one tiny sub-area amplifying light and an immediately adjacent area absorbing the generated light.

Türeci and his collaborators are now using similar ideas to pursue counterintuitive ideas for using distribution of gain and loss to make micro-lasers more efficient.

The researchers' ideas for taking advantage of loss derive from their study of mathematical constructs called "non-Hermitian" matrices in which a normally symmetric table of values becomes asymmetric. Türeci said the work is related to certain ideas of quantum physics in which the fundamental symmetries of time and space in nature can break down even though the equations used to describe the system continue to maintain perfect symmetry.

Over the past several years, Türeci and his collaborators at Vienna worked to show how the mathematical anomalies at the heart of this work, called "exceptional points," could be manifested in an actual system. In 2012 <sup>3</sup>, the team published a paper in the journal *Physical Review Letters* demonstrating computer simulations of a laser system that shuts off as energy is being added. In the current *Nature Communications* paper, the researchers created an experimental realization of their theory using a light source known as a [quantum cascade laser](#).

The researchers report in the article that results could be of particular value in creating "lab-on-a-chip" devices – instruments that pack tiny optical devices onto a single computer chip. Understanding how multiple optical devices interact could provide ways to manipulate their performance electronically in previously unforeseen ways. Taking advantage of the way loss and gain are distributed within tightly coupled

laser systems could lead to new types of highly accurate sensors, the researchers said.

"Our approach provides a whole new set of levers to create unforeseen and useful behaviors," Türeci said.

**More information:** 1. *Nature Communications* 13 June 2014. [DOI: 10.1038/ncomms5034](https://doi.org/10.1038/ncomms5034)

2. *Science* 2 May 2008. [DOI: 10.1126/science.1155311](https://doi.org/10.1126/science.1155311)

3. *Physical Review Letters* 24 April 2012. [DOI: 10.1103/PhysRevLett.108.173901](https://doi.org/10.1103/PhysRevLett.108.173901)

Provided by Princeton University

Citation: Strange physics turns off laser (2014, June 17) retrieved 24 April 2024 from <https://phys.org/news/2014-06-strange-physics-laser.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.