


Researchers develop an extremely simple procedure to obtain nanosized graphenes

July 17 2014



Atomic force microscopy (AFM) image of a clover-shaped nanographenes.
Credit: Bruno Schuler, IBM – Zurich

The prestigious journal *Angewandte Chemie* has recently published a work by CiQUS researchers (University of Santiago de Compostela, Spain) in collaboration with IBM Research – Zurich (Switzerland), which describes an extremely simple method to obtain high quality

nanographenes from easily available organic compounds.

Graphene is considered an unique material, which is leading to the emergence of a completely new technology. One of the biggest challenges in this new field is the development of methodologies for the preparation on this material with nanometric size and high quality: if the researchers get a perfect control over their size and geometry, they could explore new applications for high-performance electronic devices. The method discovered by CiQUS researchers allows to obtain well-defined nanographenes in one-pot from perylene, a very common organic compound.

This method is based on the reactivity of a group of molecules named arynes, which can act as "molecular glue" to paste graphene fragments together. The clover-shaped nanographenes obtained in this research were deposited on ultrathin insulating films, and imaged with [atomic resolution](#) by atomic force microscopy (AFM). The preparation of these materials with different size and shapes could be crucial to build [graphene](#)-based electronic circuits, molecular machinery and/or single molecule electronic devices.

The research work, led by Prof. Diego Peña, is a contribution of the research group COMMO, COMMO -belonging to CiQUS-, and includes the participation of Prof. Enrique Guitián, Prof. Dolores Pérez and the PhD student Sara Collazos. This COMMO group is pioneer on the synthesis of nanosized graphenes following bottom-up approaches through chemical methods in solution. The IBM Group ([link is external](#)) is specialist in the use of AFM with atomic resolution and in this work Dr. Leo Gross, Dr. Gerhard Meyer and PhD student Bruno Schuler were involved.

Both groups have already collaborated and published previous results in prestigious chemistry scientific journals as *Science* (2012), and currently

they take part of the Large European Project PAMS (Planar Atomic and Molecular Scale Devices) (link is external), with a total budget over 9 million Euros. PAMS main objective is to develop [electronic devices](#) of nanometric-scale size, in order to get the extreme miniaturization of the equipment used in information technology and communication.

More information: Schuler, B., Collazos, S., Gross, L., Meyer, G., Pérez, D., Guitián, E. and Peña, D. (2014), "From Perylene to a 22-Ring Aromatic Hydrocarbon in One-Pot." *Angew. Chem. Int. Ed.*. [DOI: 10.1002/anie.201403707](#)

Provided by IBM

Citation: Researchers develop an extremely simple procedure to obtain nanosized graphenes (2014, July 17) retrieved 9 April 2024 from <https://phys.org/news/2014-07-extremely-simple-procedure-nanosized-graphenes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
