

Graphene surfaces on photonic racetracks

July 28 2014, by Daniel Cochlin

In an article published in *Optics Express*, scientists from The University of Manchester describe how graphene can be wrapped around a silicon wire, or waveguide, and modify the transmission of light through it.

Just as information can be carried by electrons in metal wires in a microchip, photons of light can carry information through silicon waveguides to form a photonic microchip. Photonic microchips are often regarded as the future of computer processing and telecommunications because of the vastly increased speeds of operation and bandwidth enhancements.

Perhaps more surprisingly, they are also finding applications as highly sensitive bio-chemical sensor devices.

In this work led by University of Manchester scientists Dr Aravind Vijayaraghavan and Dr Iain Crowe in collaboration with scientists from University of Southampton, graphene coatings have been applied to what are known as 'racetrack resonators' – [waveguide](#) loops shaped like oval racetracks – to form a potentially novel device architecture.

Dr Vijayaraghavan explains: "When light travels around such a racetrack, some of the light 'leaks' out of the waveguide surface, and this 'evanescent field' can be used for chemical sensing applications. A coating of graphene on the surface of a waveguide can be used to add further capability to such a sensor, such as making it more sensitive and selective.

"In this paper, we have calculated how much light is absorbed by the graphene when it coats the waveguide, and recommend optimum conditions for the graphene coating to serve as a sensor enhancement layer."

Dr Crowe added: "The addition of the graphene layer to our [silicon waveguide](#) dramatically alters the way the light is guided through the device meaning that the light will interact even more strongly with surface deposited molecules, when employed in a sensor device.

"This strong interaction between the light and the [graphene](#) layer means that the device could also be used to improve the detection of light itself, at very low levels and across a broad range of frequencies when employed in a device known as a 'photo-detector'.

Provided by University of Manchester

Citation: Graphene surfaces on photonic racetracks (2014, July 28) retrieved 20 April 2024 from <https://phys.org/news/2014-07-graphene-surfaces-photonic-racetracks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.