

## Merging galaxies illuminate the cosmic food chain

July 1 2014, by Steve Jefferson



The Umbrella Galaxy takes its name from a mysterious feature seen on the left here, that is now found to be debris from a tiny galaxy, only a 50th its size, shredded apart by gravity. The image is a combination of data from the 0.5-meter BlackBird Remote Observatory Telescope and Suprime-Cam on the 8-meter Subaru Telescope. The inset shows a small cluster of stars embedded in the stream, which marks the center of the disrupted galaxy. Credit: R. JAY GABANY

(Phys.org) —Scientists studying a 'twin' of the Milky Way have used the W. M. Keck Observatory and Subaru Observatory to accurately model



how it is swallowing another, smaller galaxy. Their findings have opened the way to a better understanding of how structure forms in the universe and are being published in the *Monthly Notices of the Royal Astronomical Society* this week.

The work, led by Caroline Foster of the Australian Astronomical Observatory, has used the Umbrella (NGC 4651) galaxy to reveal insights in galactic behavior.

The Umbrella lies 62 million light-years away, in the northern constellation of Coma Berenices. Its faint parasol is composed of a stellar stream, thought to be the remnants of a smaller galaxy being pulled apart by the large galaxy's intense gravitational field. The Umbrella will eventually absorb this small galaxy completely.

The merging of <u>small galaxies</u> into larger ones is common throughout the universe, but because the shredded galaxies are so faint it has been hard to extract details in three-dimensions about how such mergers proceed. Using the most powerful optical facilities in the world, the twin, 10-meter Keck Observatory and the 8-meter Subaru Telescope, near the summit of Mauna Kea, Foster and her collaborators have determined enough about the character of the merger to provide a detailed model of how and when it occurred.





In this three-dimensional, rotating computer model of the Umbrella galaxy, the disk of the main galaxy is shown by blue circles. The path of the dwarf galaxy through space is shown by a green curve. The white dots show stars that once belonged to the dwarf galaxy but have now been ripped off by tidal forces into a long stream of stars. Credit: N. SINGH/UCSC

After taking panoramic images of the Umbrella with Suprime-Cam on Subaru, the scientists used the DEIMOS instrument, installed on the Keck II telescope, to map out the motions of the stream and hence determine how the galaxy is being shredded.

The stars in the stream are incredibly faint, so it was necessary to use a proxy technique to measure the speeds of brighter tracer objects moving along with the stream stars. These bright tracers include globular star clusters, planetary nebulae (dying stars that glow like neon lights), and patches of glowing hydrogen gas.

"This is important because our whole concept about what galaxies are



and how they grow has not been fully verified," said co-author Aaron Romanowsky, an astronomer at both San José State University and University of California Observatories. "We think they are constantly consuming smaller galaxies as part of a cosmic food chain, all pulled together by a mysterious form of invisible 'dark matter'. When a galaxy is torn apart, we sometimes get a glimpse of the hidden vista because the stripping process lights it up. That's what occurred here."

"Through new techniques we have been able to measure the movements of the stars in the very distant, very faint, stellar stream in the Umbrella," Foster said. "This allows us, for the first time, to reconstruct the history of the system."

"Being able to study streams this far away means that we can reconstruct the assembly histories of many more galaxies," Romanowsky said. "In turn that means we can get a handle on how often these 'minor mergers'—thought to be an important way that galaxies grow—actually occur. We can also map out the orbits of the stellar streams to test the pull of gravity for exotic effects, much like the Moon going around the Earth but without having to wait 300 million years for the orbit to complete."

The present work is a follow-up to a 2010 study, led by Dr. David Martínez-Delgado (University of Heidelberg), which used small robotic telescopes to image eight isolated spiral galaxies, and found the signs of mergers—shells, clouds and arcs of tidal debris—in six of them.

The W. M. Keck Observatory operates the largest, most scientifically productive telescopes on Earth. The two, 10-meter optical/infrared telescopes on the summit of Mauna Kea on the Island of Hawaii feature a suite of advanced instruments including imagers, multi-object spectrographs, high-resolution spectrographs, integral-field spectroscopy and world-leading laser guide star adaptive optics systems.



DEIMOS (the DEep Imaging and Multi-Object Spectrograph) boasts the largest field of view (16.7 arcmin by 5 arcmin) of any of the Keck instruments, and the largest number of pixels (64 Mpix). It is used primarily in its multi-object mode, obtaining simultaneous spectra of up to 130 galaxies or stars. Astronomers study fields of distant <u>galaxies</u> with DEIMOS, efficiently probing the most distant corners of the universe with high sensitivity.

Provided by W. M. Keck Observatory

Citation: Merging galaxies illuminate the cosmic food chain (2014, July 1) retrieved 3 May 2024 from <u>https://phys.org/news/2014-07-merging-galaxies-illuminate-cosmic-food.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.