

Social network research may boost prairie dog conservation efforts

July 28 2014, by Matt Shipman



Credit: Jennifer Verdolin

Researchers using statistical tools to map social connections in prairie dogs have uncovered relationships that escaped traditional observational techniques, shedding light on prairie dog communities that may help



limit the spread of bubonic plague and guide future conservation efforts. The work was done by researchers from North Carolina State University and the National Evolutionary Synthesis Center (NESCent).

"Prairie dogs are increasingly rare and are subject to <u>bubonic plague</u>," says Dr. Jennifer Verdolin, lead author of a paper on the work and an animal behavior researcher at NESCent. "This work improves our understanding of the social dynamics within a prairie dog colony, which could help us find ways to mitigate the spread of plague. It also helps us understand how these <u>social groups</u> are connected, so we can limit the disruption to prairie dog populations when we need to relocate them for conservation purposes."

"This project also emphasizes how useful social <u>network theory</u> can be when applied to population biology," says Amanda Traud, co-author of the paper and a Ph.D. candidate in biomathematics at NC State. "And we also developed new tools for this project that can be applied elsewhere and help us use social network theory more effectively to study other species.

"For example, we developed new code that can identify individuals that serve as 'hubs' or 'bridges,' connecting different groups within a population," Traud says.

The researchers used data Verdolin collected on three Gunnison's prairie dog (Cynomys gunnisoni) colonies in Arizona, ranging in size from 60 to 200 individuals. Each colony is made up of numerous, smaller social groups that can consist of up to 15 individuals.

Traud developed statistical tools using social network theory to analyze one year's worth of data on the three colonies, focusing specifically on "greet kissing," in which prairie dogs briefly lock teeth with each other. If individuals are from the same social group, they part amicably. If they



are from different groups, they usually fight or chase one another after greet kissing.

By looking at data on greet kissing, Traud established which <u>prairie dogs</u> were in the same group. These groups were consistent with the groups Verdolin had determined, using conventional observational methods of behavior and location.

However, the math-driven social network theory approach also identified substructures within those social groups and key "bridge" individuals that connected different groups – findings that may have significant conservation applications.

"For example, there's a possibility that we could slow or stop the spread of plague in a colony by relocating these bridge individuals," Traud says.

"By the same token," Verdolin says, "conservationists could use this data to be sure to relocate all of a relevant social group, rather than splitting up groups. That could improve the group's chances of thriving in a new environment."

More information: Jennifer L. Verdolin, Amanda L. Traud, Robert R. Dunn, "Key players and hierarchical organization of prairie dog social networks," *Ecological Complexity*, Volume 19, September 2014, Pages 140-147, ISSN 1476-945X, DOI: 10.1016/j.ecocom.2014.06.003.

Provided by North Carolina State University

Citation: Social network research may boost prairie dog conservation efforts (2014, July 28) retrieved 6 May 2024 from <u>https://phys.org/news/2014-07-social-network-boost-prairie-dog.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.