

And then there were 10—unexpected diversity in New Zealand kanuka genus *Kunzea*

August 26 2014



A *Kunzea robusta* flowering branchlet, in Waitakere Ranges, West Auckland, North Island New Zealand. Credit: Peter J. de Lange

At the stroke of a pen a New Zealand endemic tree has for the last 31 years been incorrectly regarded the same as a group of 'weedy'

Australian shrubs and small trees. A New Zealand botanist has completed a 15-year study to reveal some surprises and discover astonishing cryptic diversity behind what was long considered a single tree species. The study was published in the open access journal *PhytoKeys*.

Known to botanists as *Kunzea ericoides*, this species was one of the many discoveries made in the north-western South Island of New Zealand by Jules Sébastien César Dumont d'Urville during the first voyage of the *Astrolabe* (1826-1829).

Initially described by French Professor of botany Achille Richard as *Leptospermum ericoides*, this species and the allied New Zealand endemic *Leptospermum sinclairii* were merged in 1983 with three other Australian species under the oldest available name (*L. ericoides*) as a new combination in the related genus *Kunzea*.

The saying that 'bad taxonomy kills' could not be more true, as the decision to merge the Australian and New Zealand species into the one shared species – *Kunzea ericoides* – resulted in absolute mayhem as a range of important New Zealand trees and shrubs were now regarded as identical to their distant Australian relatives all of which are serious agricultural pests known there as Bursaria.

The present study draws on data obtained from experimental hybridisations, cytological investigations, modern molecular analyses and ecological studies published elsewhere by the author of this revision Dr Peter J. de Lange. The *PhytoKeys* treatment by de Lange concludes that *Kunzea ericoides* is not shared between Australia and New Zealand; confirms the distinctiveness of the narrow-range rhyolite rock endemic *K. sinclairii*; and elevates another much misunderstood variety to species rank as *K. linearis*.

De Lange also recognises further seven species new to science. One, *Kunzea robusta* is actually the most common *Kunzea* in New Zealand which despite being first recognised as distinct by the missionary botanist William Colenso in the 1840s, was denied formal recognition for some 170 years until now. Aside from *Kunzea robusta* de Lange recognises two other common new tree species (*K. amathicola* and *K. serotina*), and another four narrow-range endemics - *K. salterae*, *K. tenuicaulis*, *K. toelkenii*, and *K. triregensis*. One of these species, *K. tenuicaulis* is known only from active geothermal fields.



A *Kunzea ericoides* flowering branchlet, in Pupu Springs, Golden Bay, North-west Nelson, South Island New Zealand. Credit: Peter J. de Lange

Aside from the recognition of 10 endemic New Zealand species from what had been regarded as one in 1983, de Lange also reveals that the indigenous inhabitants of New Zealand the Maori already knew of many of these species, which they recognised by names which referred to their growth habit or wood properties.

It is shown that far from being serious agricultural weeds all the New Zealand *Kunzea* are important keystone species, either forming their own distinct forest types or helping to heal damaged forest. Furthermore the New Zealand *Kunzea* species provide an important habitat for a wealth of endemic geckos, orchids and fungi – a far cry indeed from their past much undeserved 'weed' status.

"This paper is also a brilliant showcase of how useful is the brand new *PhytoKeys* publishing platform for the advancement of taxonomy. Once published taxonomic treatments of new species and any nomenclatural changes are automatically available in advanced open access, easy to find, and are stored in various data repositories in reusable format. Such wide availability of data helps for mistakes and misunderstanding regarding previous confusions in the taxonomy of a certain [species](#) to be avoided in future," comments Prof. Lyubomir Penev, managing director of [Pensoft Publishers](#).

More information: de Lange PJ (2014) A revision of the New Zealand *Kunzea ericoides* (Myrtaceae) complex. *PhytoKeys* 40: 1-185. [DOI: 10.3897/phytokeys.40.7973](https://doi.org/10.3897/phytokeys.40.7973)

Provided by Pensoft Publishers

Citation: And then there were 10—unexpected diversity in New Zealand kanuka genus *Kunzea* (2014, August 26) retrieved 23 May 2024 from <https://phys.org/news/2014-08-10unexpected->

diversity-zealand-kanuka-genus.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.