

## Acid ions are more than spectators

August 7 2014, by David Bradley



The experimental prediction and theoretical confirmation of persistent ion pairing between hydronium and chloride provides a unified description of the molecular structure of concentrated hydrogen-chloride solutions. The partial decompositions into Cl–OH3O+ and Cl–OH2O are shown in orange and green,



respectively, with the structural portions depicted as insets in (b) (water hydrogens and oxygens are rendered as small and large gray-scaled spheres, respectively. Charge is indicated with "+" or "-", and the choride is rendered as a green sphere). (d) is the experimental set-up.

(Phys.org) —X-ray absorption fine structure (EXAFS) measurements carried out at the U.S. Department of Energy's (DOE's) Advanced Photon Source, coupled with state-of-the-art density functional theory (DFT) simulations reveal that strong acids, such as hydrochloric acid, form counter-ion pairs in solution across all concentration, a result that had not been seen in gas-phase studies. The discovery suggests that it is not simply the release of protons—hydrogen ions—that is important for the properties of acids.

When a strong <u>acid</u>, such as hydrochloric acid, forms a solution in water, the <u>hydrogen ions</u> break away from the compound and give rise to the acidity of the solution. These hydrogen ions can diffuse quickly in the solution and attack other chemicals present, neutralizing basic, or alkaline, substances, or corroding metals and reacting with organic substances. Meanwhile, the counter ion (the chloride in the example of hydrochloric acid) was assumed to form independent, negatively charged solvated fragments, but little was known about the properties of the counter ions or precisely what ion fragments they generated.

Chemists had computer models at their disposal, often based on gasphase studies of the particles involved. However, acids are present in the condensed phase and, thus, the role of the condensed phase environment, as opposed to an isolated molecule in the vapor, should be considered. Moreover, as the researchers in this study report in a cover article for the *Journal of Physical Chemistry B*, the gas-phase studies are not representative of the more common acidic solution.



Countless chemical reactions hinge on the presence of an excess of protons in the reaction solution, formed by the dissociation of a strong acid into dissolved hydrogen ions and the counter ion. Indeed, these so-called "BrØnsted acids" are used in many industrially important chemical reactions, are involved in numerous biological reactions, and play a role in atmospheric chemistry.

Chemists have assumed that the dissociation of a strong acid has two steps. First, the hydrogen-bearing, or protic, acid bonds loosely to a water molecule and the electrical charge is redistributed so that the counter ion becomes more negative while the water molecule gains a positive charge. In the second step, the counter ion is released as a freefloating chloride ion, for instance, and the positively charged water molecule grabs the proton from the acid to form H3O+, the hydronium ion. The extra proton on this chemical species is free to diffuse through the solution, endowing the solution with its acidic character. These researchers, from Pacific Northwest National Laboratory and Argonne National Laboratory, have provided a molecular picture of deviations from ideal acid dissociation.

The team carried out extended x-ray absorption fine structure studies at the X-ray Science Division (XSD) 20-BM beamline at the Argonne Advanced Photon Source, a DOE Office of Science User Facility. They combined these data with molecular dynamics and corroborated earlier neutron and x-ray diffraction data. Specifically, the bond distance between chloride ion and the oxygen in the hydronium ion comprising the contact ion pair significantly shorter than the interaction between chloride and the oxygen of water in the ideal dissociation picture. This difference in distance and geometry can be measured. Furthermore, the newly discovered molecular moieties can be used to re-interpret in a new light long-standing neutron and x-ray diffraction data.

Thus, the contact ion pair between the H3O+ and chloride suggests that



the long-misunderstood counter ion of a protic acid is not a mere spectator but is integral to defining the properties of acids across a broad concentration range. Indeed, across the concentration range, the team demonstrates that the contact <u>ion</u> pair is ubiquitous with very few free protons in solution at the highest concentrations. The study provides a template to investigate and understand the details of other acids and their chemistries under a variety of different environments including interfaces.

**More information:** Marcel D. Baer1, John L. Fulton1, Mahalingam Balasubramanian2, Gregory K. Schenter1, and Christopher J. Mundy1\*, "Persistent Ion Pairing in Aqueous Hydrochloric Acid," *J. Phys. Chem. B* 118, 7211 (2014). DOI: 10.1021/jp501091h

Provided by Argonne National Laboratory

Citation: Acid ions are more than spectators (2014, August 7) retrieved 23 June 2024 from <u>https://phys.org/news/2014-08-acid-ions-spectators.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.