

Air quality in San Joaquin Valley improving according to study

August 21 2014, by Lynn Yarris



Air quality in California's San Joaquin Valley has long ranked as one of the worst in the nation but NO_x emission controls are improving the situation.

(Phys.org) —Air quality in California's San Joaquin Valley, which for years has been ranked as one of the worst in the United States has improved over the past decade and looks to improve even more in the future. Speaking at the national meeting of the American Chemical Society (ACS) in San Francisco, chemist Sally Pusede described how efforts to reduce nitrogen oxide (NO_x) emissions have dramatically improved air quality in the valley over the past decade and should be

even more effective over the next ten years. Pusede is an alumna of the research group of Ronald Cohen, a faculty scientist with the Lawrence Berkeley National Laboratory (Berkeley Lab) and the University of California (UC) Berkeley.

"Our study showed that NO_x reductions in the San Joaquin Valley as the result of implemented controls have dramatically decreased the very large fraction of total aerosol that is ammonium nitrate," Pusede said. "The good news is that in the next ten years, the additional NO_x reductions that are expected from new controls on heavy-duty diesel trucks will halve the frequency of days in the Valley on which the air quality standard is exceeded."

According to the American Lung Association, the San Joaquin Valley experiences the worst pollution in the U.S. by small particulate matter (PM_{2.5}) – particles of soot, dust, unburned fuel, and matter chemically formed in the atmosphere that are 2.5 micrometers or less in size. While PM_{2.5} is a wintertime problem, in the summer the San Joaquin Valley is second only to Los Angeles for having the worst ozone pollution levels.

"As a result, extremely poor air quality in the San Joaquin Valley is a year round problem," Pusede said.

Pusede and Cohen, a leading authority on atmospheric chemistry with Berkeley Lab's Environmental Energy Technologies Division, examined 15 years of data on NO_x and other emissions, and their chemical products in three key San Joaquin Valley cities – Fresno, Visalia and Bakersfield.

"NO_x abundances have fallen steadily over the last fifteen years and patterns in anthropogenic sources result in two-fold lower NO_x emissions on weekends than weekdays largely without co-occurring changes in other emissions," Pusede said. "These trends taken together

provide a useful constraint on the NO_x dependence of the driving chemical mechanisms of both ozone and aerosol."

Pusede and Cohen used this NO_x constraint to interpret trends in wintertime PM_{2.5} over the last decade in San Joaquin Valley where a large portion of the total aerosol mass is [ammonium nitrate](#). They also looked at the impact on the formation of ozone, which is a product of nitrogen oxide chemistry.

"We found that the NO_x controls that have reduced PM_{2.5} levels are also poised to be incredibly effective at decreasing the frequency of days with high ozone in the region," Pusede said.

Added Cohen in commenting on the research, "The bottom line is that [air quality](#) in the San Joaquin valley is substantially better as a result of NO_x reductions and we predict it will get better still as new regulations take effect."

Provided by Lawrence Berkeley National Laboratory

Citation: Air quality in San Joaquin Valley improving according to study (2014, August 21)
retrieved 23 April 2024 from <https://phys.org/news/2014-08-air-quality-san-joaquin-valley.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--