

Alternate mechanism of species formation picks up support, thanks to a South American ant

August 21 2014



A queen ant of the host species *Mycocepurus goeldii*. Credit: Scott Solomon

A newly-discovered species of ant supports a controversial theory of species formation. The ant, known to live only under a single eucalyptus tree on the São Paulo State University campus in Brazil, branched off

from its original species while living in the same colony, something thought rare in current models of evolutionary development.

"Most new species come about in geographic isolation," said Christian Rabeling, assistant professor of biology at the University of Rochester. "We now have evidence that speciation can take place within a single colony."

The findings by Rabeling and the research team were published today in the journal *Current Biology*.

In discovering the parasitic *Mycocepurus castrator*, Rabeling and his colleagues uncovered an example of a still-controversial theory known as sympatric speciation, which occurs when a new species develops while sharing the same geographic area with its parent species, yet reproducing on its own. "While sympatric speciation is more difficult to prove," said Rabeling, "we believe we are in the process of actually documenting a particular kind of evolution-in-progress."

New species are formed when its members are no longer able to reproduce with members of the parent species. The commonly-accepted mechanism is called allopatric speciation, in which geographic barriers—such as mountains—separate members of a group, causing them to evolve independently.



A queen ant of the parasitic species *Mycocepurus castrator*. Credit: Scott Solomon

"Since Darwin's *Origin of Species*, evolutionary biologists have long debated whether two species can evolve from a common ancestor without being geographically isolated from each other," said Ted Schultz, curator of ants at the Smithsonian's National Museum of Natural History and co-author of the study. "With this study, we offer a compelling case for sympatric evolution that will open new conversations in the debate about speciation in these ants, social insects and evolutionary biology more generally." *M. castrator* is not simply another ant in the colony; it's a parasite that lives with—and off of—its [host](#), *Mycocepurus goeldii*. The host is a fungus-growing ant that cultivates fungus for its nutritional value, both for itself and, indirectly, for its parasite, which does not participate in the work of growing the

fungus garden. That led the researchers to study the genetic relationships of all fungus-growing ants in South America, including all 11 known species of the genus *Mycocepurus*, to determine whether the parasite did evolve from its presumed host. They found that the parasitic ants were, indeed, genetically very close to *M. goeldii*, but not to the other ant species.

They also determined that the parasitic ants were no longer reproductively compatible with the host ants—making them a unique species—and had stopped reproducing with their host a mere 37,000 years ago—a very short period on the evolutionary scale.

A big clue for the research team was found by comparing the ants' genes, both in the cell's nucleus as well as in the mitochondria—the energy-producing structures in the cells. Genes are made of units called nucleotides, and Rabeling found that the sequencing of those nucleotides in the mitochondria is beginning to look different from what is found in the host ants, but that the genes in the nucleus still have traces of the relationship between host and parasite, leading him to conclude that *M. castrator* has begun to evolve away from its host.

Rabeling explained that just comparing some nuclear and mitochondrial genes may not be enough to demonstrate that the parasitic ants are a completely new species. "We are now sequencing the entire mitochondrial and nuclear genomes of these parasitic ants and their host in an effort to confirm speciation."

The parasitic ants need to exercise discretion because taking advantage of the host species is considered taboo in ant society. Offending ants have been known to be killed by worker mobs. As a result, the parasitic queen of the [new species](#) has evolved into a smaller size, making them difficult to distinguish from a host worker.

Host queens and males reproduce in an aerial ceremony only during a particular season when it begins to rain. Rabeling found that the parasitic queens and males, needing to be more discreet about their reproductive activities, ignore seasonal cues. By needing to hide their parasitic identity, *M. castrator* males and females lost their special adaptations that allowed them to reproduce in flight, making it impossible for them to sexually interact with their host [species](#).

Provided by University of Rochester

Citation: Alternate mechanism of species formation picks up support, thanks to a South American ant (2014, August 21) retrieved 2 May 2024 from <https://phys.org/news/2014-08-alternate-mechanism-species-formation-south.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--