

Big snowstorms will still occur in Northern Hemisphere following global warming, study finds

August 27 2014, by Jennifer Chu



Coated in ice, power and telephone lines sag and often break, resulting in power outages. Credit: Public Domain

If ever there were a silver lining to global warming, it might be the prospect of milder winters. After all, it stands to reason that a warmer climate would generate less snow.



But a new MIT study suggests that you shouldn't put your shovels away just yet. While most areas in the Northern Hemisphere will likely experience less <u>snowfall</u> throughout a season, the study concludes that extreme snow events will still occur, even in a future with significant warming. That means that, for example, places like Boston may see less snowy winters overall, punctuated in some years by blizzards that drop a foot or two of snow.

"Many studies have looked at average snowfall over a season in climate models, but there's less known about these very heavy snowfalls," says study author Paul O'Gorman, an associate professor in MIT's Department of Earth, Atmospheric and Planetary Sciences. "In some regions, it is possible for average snowfall to decrease, but the snowfall extremes actually intensify."

O'Gorman studied daily snowfall across the Northern Hemisphere using 20 different climate models, each of which projected <u>climate change</u> over a 100-year period, given certain levels of greenhouse gas emissions. He looked at both average seasonal snowfall and extreme snowfall events under current climate conditions, and also following projected future warming.

Not surprisingly, O'Gorman found that under relatively high warming scenarios, low-elevation regions with winter temperatures initially just below freezing experienced about a 65 percent reduction in average winter snowfall. However, in these same regions, the heaviest snowstorms became only 8 percent less intense. In some higher-latitude regions, extreme snow events became more intense, depositing 10 percent more snow, even under scenarios of relatively high global warming.

"You might expect with a warmer climate there should be major changes in snowfall in general," O'Gorman says. "But that seems to be true to a



greater extent for average snowfall than for the intensities of the heaviest snowfall events."

O'Gorman has published the results of his study this week in the journal *Nature*.

Daily snowfall

For the most part, researchers have only been able to analyze snowfall on a seasonal scale, estimating a winter's average snow amounts with climate change. Such analyses, while useful, only paint a broad picture of snowfall's response to global warming, and may miss specific events, like a large blizzard that may occur over a day or two.

Daily snowfall in a range of climate model simulations has recently been made available through the Coupled Model Intercomparison Project—a growing archive of climate modeling output, including snowfall, that modeling centers and researchers around the world contribute to and analyze.

O'Gorman analyzed daily snow amounts from simulations with 20 different climate models in the archive. Each model simulated a "control climate," for the years 1981 to 2000, as well as a "warm climate," for the years 2081 to 2100, assuming relatively high emissions of greenhouse gases.

Over this 100-year period, O'Gorman found that average snowfall decreased substantially in many Northern Hemisphere regions in warmclimate scenarios compared with the milder control climates, but that snowfall amounts in the largest snowstorms did not decrease to the same extent.

He warned, however, that changes in snowfall extremes can be larger in



regions with little snowfall to begin with, such as the southwestern United States. He also notes that while this study focuses on percentage changes in the amount of snowfall in extreme snowfall events, there can be larger changes in the frequency of such events.

From the simulations, O'Gorman found that it takes greater climate warming to reduce the intensity of extreme snowstorms than to reduce average seasonal snowfall. Specifically, a region would experience less seasonal snow if average winter temperatures were initially above minus 14 degrees Celsius (7 degrees Fahrenheit). But the heaviest snowstorms would become less intense only above minus 9 C (16 F).

A sweet spot for extreme snowfall

What's more, O'Gorman found that there's a narrow daily temperature range, just below the freezing point, in which extreme snow events tend to occur—a sweet spot that does not change with <u>global warming</u>. This is in contrast to average snow events, which may occur over a broader temperature range.

"People may know the expression, 'It's too cold to snow'—if it's very cold, there is too little water vapor in the air to support a very heavy snowfall, and if it's too warm, most of the precipitation will fall as rain," O'Gorman says. "Snowfall extremes still occur in the same narrow temperature range with climate change, and so they respond differently to <u>climate</u> change compared to rainfall extremes or average snowfall."

More information: Contrasting responses of mean and extreme snowfall to climate change, *Nature*, <u>dx.doi.org/10.1038/nature13625</u>

Provided by Massachusetts Institute of Technology



Citation: Big snowstorms will still occur in Northern Hemisphere following global warming, study finds (2014, August 27) retrieved 1 May 2024 from <u>https://phys.org/news/2014-08-big-snowstorms-northern-hemisphere-global.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.