

# Canola flowers faster with heat genes

August 21 2014, by David Stacey

---



(Phys.org) —A problem that has puzzled canola breeders for years has been solved by researchers from The University of Western Australia - and the results could provide a vital breakthrough in understanding the impact of increasing global temperatures on crop flowering.

The key to understanding what makes Australian [canola](#) flower earlier than its Canadian and European counterparts lies in the [genes](#).

Associate Professor Matthew Nelson from UWA's Institute of Agriculture and School of Plant Biology has identified that [heat](#)-responsive genes are responsible for [flowering time](#) in Australian spring-type and European summer-type canola. This is the first time such genes have been reported to influence flowering time in canola.

Australian canola is quite distinct from its Canadian and European counterparts - it flowers much earlier. Plant breeders cannot simply transfer varieties from Canada or Europe into Australia as they flower much too late for the Australian environment.

"We took a European summer-type canola, crossed it with Monty, a typical early flowering Australian variety, and analysed the progeny for variation in flowering time," Associate Professor Nelson said.

"There was a huge variation from about 30 days to 160 days in our typical Australian environment. This was totally unexpected and we showed there are several forms of these heat-responsive genes controlling flowering time."

The research indicated that the European plants required much more accumulated heat (thermal time) to flower than the Australian plants.

"Until now, most researchers assumed that long summer days in Europe and Canada triggered flowering, not heat," Associate Professor Nelson said. "Now we know that long days are only a minor part of the story."

"Understanding this complex process is important as breeders alter the adaptation of crops to a new and changing environment," research team leader Winthrop Professor Wallace Cowling said. "International canola breeders will use this information to re-establish the correct flowering time in canola when they cross between Australian types and summer annual types in the northern hemisphere.

"With increasing [global temperatures](#), or in low rainfall environments, it will be possible to 'mix and match' forms of these heat genes to achieve the target flowering date."

Since day-length affected both European and Australian varieties

equally, the research group can now focus on heat genes to predict the outcomes from crossing European or Canadian canola with Australian canola and explain what caused these genes to respond to accumulated heat.

**More information:** The paper: Quantitative trait loci for thermal time to flowering and photoperiod responsiveness discovered in summer annual-type *Brassica napus* L. is available online:

[www.plosone.org/article/fetchO...ject.action?uri=info](http://www.plosone.org/article/fetchO...ject.action?uri=info)

[%3Adoi%2F10.1371%2Fjournal.pone.0102611&representation=PDF](http://doi.org/10.1371/journal.pone.0102611&representation=PDF)

Provided by University of Western Australia

Citation: Canola flowers faster with heat genes (2014, August 21) retrieved 23 April 2024 from <https://phys.org/news/2014-08-canola-faster-genes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.