


# Cassini tracks clouds developing over a Titan sea

August 12 2014, by Preston Dyches

---




As NASA's Cassini spacecraft sped away from Titan following a relatively close flyby, its cameras monitored the moon's northern polar region, capturing signs of renewed cloud activity. Credit: NASA

(Phys.org) —NASA's Cassini spacecraft recently captured images of clouds moving across the northern hydrocarbon seas of Saturn's moon Titan. This renewed weather activity, considered overdue by researchers, could finally signal the onset of summer storms that atmospheric models have long predicted.

A movie showing the clouds' movement can be seen below.

The Cassini spacecraft obtained the new views in late July, as it receded from Titan after a close flyby. Cassini tracked the system of clouds developing and dissipating over the large methane sea known as Ligeia Mare for more than two days. Measurements of cloud motions indicate wind speeds of around 7 to 10 mph (3 to 4.5 meters per second).

For several years after Cassini's 2004 arrival in the Saturn system, scientists frequently observed cloud activity near Titan's south pole, which was experiencing late summer at the time. Clouds continued to be observed as spring came to Titan's northern hemisphere. But since a huge storm swept across the icy moon's low latitudes in late 2010, only a few small clouds have been observed anywhere on the icy moon. The lack of [cloud activity](#) has surprised researchers, as computer simulations of Titan's atmospheric circulation predicted that clouds would increase in the north as summer approached, bringing increasingly warm temperatures to the atmosphere there.



This animated sequence of Cassini images shows methane clouds moving above the large methane sea on Saturn's moon Titan known as Ligeia Mare. The spacecraft captured the views between July 20 and July 22, 2014, as it departed Titan following a flyby. Cassini tracked the system of clouds as it developed and dissipated over Ligeia Mare during this two-day period. Measurements of the cloud motions indicate wind speeds of around 7 to 10 miles per hour (3 to 4.5 meters per second). The timing between exposures in the sequence varies. In particular, there is a 17.5-hour jump between the second and third frames. Most other frames are separated by one to two hours. Credit: NASA

"We're eager to find out if the clouds' appearance signals the beginning

of summer weather patterns, or if it is an isolated occurrence," said Elizabeth Turtle, a Cassini imaging team associate at the Johns Hopkins University Applied Physics Lab in Laurel, Maryland. "Also, how are the [clouds](#) related to the seas? Did Cassini just happen catch them over the seas, or do they form there preferentially?"

A year on Titan lasts about 30 Earth years, with each season lasting about seven years. Observing seasonal changes on Titan will continue to be a major goal for the Cassini mission as summer comes to Titan's north and the southern latitudes fall into winter darkness.

Provided by NASA

Citation: Cassini tracks clouds developing over a Titan sea (2014, August 12) retrieved 22 June 2024 from <https://phys.org/news/2014-08-cassini-tracks-clouds-titan-sea.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.