


Copper foam turns carbon dioxide into useful chemicals

August 12 2014



Scientists at Brown University's Center for Capture and Conversion of CO₂ have discovered that copper foam could provide a new way of converting excess CO₂ into useful industrial chemicals. The foam offers sponge-like pores and channels, providing more active sites for CO₂ reactions than a simple surface. Credit: Palmore Lab / Brown University

A catalyst made from a foamy form of copper has vastly different electrochemical properties from catalysts made with smooth copper in

reactions involving carbon dioxide, a new study shows. The research, by scientists in Brown University's Center for the Capture and Conversion of CO₂, suggests that copper foams could provide a new way of converting excess CO₂ into useful industrial chemicals.

The research is published in the journal *ACS Catalysis*.

As levels of [carbon dioxide](#) in the atmosphere continue to rise, researchers are looking for ways to make use of it. One approach is to capture CO₂ emitted from power plants and other facilities and use it as a carbon source to make [industrial chemicals](#), most of which are currently made from fossil fuels. The problem is that CO₂ is extremely stable, and reducing it to a reactive and useful form isn't easy.

"Copper has been studied for a long time as an electrocatalyst for CO₂ reduction, and it's the only metal shown to be able to reduce CO₂ to useful hydrocarbons," said Tayhas Palmore, professor of engineering and senior author of the new research. "There was some indication that if you roughen the surface of planar [copper](#), it would create more active sites for reactions with CO₂."

Copper foam, which has been developed only in the last few years, provided the surface roughness that Palmore and her colleagues were looking for. The foams are made by depositing copper on a surface in the presence of hydrogen and a strong electric current. Hydrogen bubbles cause the copper to be deposited in an arrangement of sponge-like pores and channels of varying sizes.

After depositing copper foams on an electrode, the researchers set up experiments to see what kinds of products would be produced in an electrochemical reaction with CO₂ in water. The experiments were performed by Sujat Sen and Dan Liu, graduate students in chemistry working in Palmore's lab at Brown's School of Engineering.

The experiments showed that the copper foam converted CO₂ into formic acid—a compound often used as a feedstock for microbes that produce biofuels—at a much greater efficiency than planar copper. The reaction also produced small amounts of propylene, a useful hydrocarbon that's never been reported before in reactions involving copper.

"The product distribution was unique and very different from what had been reported with planar electrodes, which was a surprise," Palmore said. "We've identified another parameter to consider in the electroreduction of CO₂. It's not just the kind of metal that's responsible for the direction this chemistry goes, but also the architecture of the catalyst."

Now that it's clear that architecture matters, Palmore and her colleagues are working to see what happens when that architecture is tweaked. It's likely, she says, that pores of different depths or diameters will produce different compounds from a CO₂ feedstock. Ultimately, it might be possible to tune the copper foam toward a specific desired compound.

Palmore said she's amazed by the fact that there's still more to be learned about copper.

"People have studied electrocatalysis with copper for a couple decades now," she said. "It's remarkable that we can still make alterations to it that affect what's produced."

The work in the study is part of a larger effort by Brown's Center for the Capture and Conversion of CO₂. The Center, funded by the National Science Foundation, is exploring a variety of catalysts that can convert CO₂ into usable forms of carbon.

"The goal is to find ways to produce some of the world's largest-volume

chemicals from a sustainable [carbon source](#) that the Earth not only has in excess but urgently needs to reduce," said Palmore, who leads the center. "This is a way for us as scientists to begin thinking of how we produce industrial chemicals in more sustainable ways and control costs at the same time. The cost of commodity chemicals is going nowhere but up as long as production is dependent on [fossil fuels](#)."

Provided by Brown University

Citation: Copper foam turns carbon dioxide into useful chemicals (2014, August 12) retrieved 25 April 2024 from <https://phys.org/news/2014-08-copper-foam-carbon-dioxide-chemicals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.