

Asian monsoon much older than previously thought

September 14 2014



These 35-million-year-old snail fossils in Myanmar contain the record of the past climate in their shells. Analyses by University of Arizona geoscientist Alexis Licht and his colleagues show these snails lived in a monsoon climate that had dry winters and very rainy summers. These freshwater gastropods from the Yaw formation (37-34 million years ago) were sampled near the town of Kalewa in central Myanmar. Credit: Alexis Licht 2012

The Asian monsoon already existed 40 million years ago during a period of high atmospheric carbon dioxide and warmer temperatures, reports an international research team led by a University of Arizona geoscientist.

Scientists thought the climate pattern known as the Asian monsoon began 22-25 million years ago as a result of the uplift of the Tibetan Plateau and the Himalaya Mountains.

"It is surprising," said lead author Alexis Licht, now a research associate in the UA department of geosciences. "People thought the monsoon started much later."

The monsoon, the largest climate system in the world, governs the climate in much of mainland Asia, bringing torrential summer rains and dry winters.

Co-author Jay Quade, a UA professor of geosciences, said, "This research compellingly shows that a strong Asian monsoon system was in place at least by 35-40 million years ago."

The research by Licht and his colleagues shows the earlier start of the monsoon occurred at a time when atmospheric CO₂ was three to four times greater than it is now. The monsoon then weakened 34 million years ago when atmospheric CO₂ then decreased by 50 percent and an ice age occurred.

Licht said the study is the first to show the rise of the monsoon is as much a result of global climate as it is a result of topography. The team's paper is scheduled for early online publication in the journal *Nature* on Sept. 14.

"This finding has major consequences for the ongoing global warming," he said. "It suggests increasing the atmospheric CO₂ will increase the monsoonal precipitation significantly."



University of Arizona geoscientist Alexis Licht (bottom left) and his colleagues from the French-Burmese Paleontological Team led by Jean-Jacques Jaeger of the University of Poitiers, France (center with hiking staff) used fossils they collected in Myanmar to figure out that the Asian monsoon started at least 40 million years ago. Credit: French-Burmese Paleontological Team 2012

Unraveling the monsoon's origins required contributions from three different teams of scientists that were independently studying the environment of 40 million years ago.

All three investigations showed the monsoon climate pattern occurred 15 million years earlier than previously thought. Combining different lines of evidence from different places strengthened the group's confidence in the finding, Licht said. The climate modeling team also linked the development of the monsoon to the increased CO₂ of the time.

Licht and his colleagues at Poitiers and Nancy universities in France examined snail and mammal fossils in Myanmar. The group led by G. Dupont-Nivet and colleagues at Utrecht University in the Netherlands studied lake deposits in Xining Basin in central China. J.-B. Ladant and Y. Donnadieu of the Laboratory of Sciences of the Climate and Environment (LSCE) in Gif-sur-Yvette, France, created [climate simulations](#) of the Asian climate 40 million years ago.

A complete list of authors of the group's publication, "Asian monsoons in a late Eocene greenhouse world," is at the bottom of this release, as is a list of funding sources.

Licht didn't set out to study the origin of the monsoon.

He chose his study site in Myanmar because the area was rich in mammal fossils, including some of the earliest ancestors of modern monkeys and apes. The research, part of his doctoral work at the University of Poitiers, focused on understanding the environments those early primates inhabited. Scientists thought those primates had a habitat like the current evergreen tropical rain forests of Borneo, which do not have pronounced differences between wet and dry seasons.

To learn about the past environment, Licht analyzed 40-million-year-old freshwater snail shells and teeth of mammals to see what types of oxygen they contained. The ratio of two different forms of oxygen, oxygen-18 and oxygen-16, shows whether the animal lived in a relatively wet climate or an arid one.



Mana Rugbumrung, a researcher at the Department of Mineral Resources in Bangkok, examines the freshly excavated skull of an anthracothere, a hippopotamus-like mammal that lived 40 million years ago in what is now Myanmar. Analyses of the teeth of these ancient mammals and others from the area revealed that the animals lived in a monsoon climate that had dry winters and very rainy summers. Credit: Alexis Licht 2012

"One of the goals of the study was to document the pre-monsoonal conditions, but what we found were monsoonal conditions," he said.

To his surprise, the oxygen ratios told an unexpected story: The region had a seasonal pattern very much like the current monsoon – dry winters and very rainy summers.

"The early primates of Myanmar lived under intense seasonal stress – aridity and then monsoons," he said. "That was completely unexpected."

The team of researchers working in China found another line of evidence pointing to the existence of the monsoon about 40 million years ago. The monsoon climate pattern generates winter winds that blow dust from central Asia and deposits it in thick piles in China. The researchers found deposits of such dust dating back 41 million years ago, indicating the monsoon had occurred that long ago.

The third team's [climate](#) simulations indicated strong Asian monsoons 40 million years ago. The simulations showed the level of atmospheric CO₂ was connected to the strength of the monsoon, which was stronger 40 million years ago when CO₂ levels were higher and weakened 34 million years ago when CO₂ levels dropped.

Licht's next step is to investigate how geologically short-term increases of atmospheric CO₂ known as hyperthermals affected the monsoon's behavior 40 million years ago.

"The response of the [monsoon](#) to those hyperthermals could provide interesting analogs to the ongoing global warming," he said.

More information: Asian monsoons in a late Eocene greenhouse world, *Nature*, [DOI: 10.1038/nature13704](https://doi.org/10.1038/nature13704)

Provided by University of Arizona

Citation: Asian monsoon much older than previously thought (2014, September 14) retrieved 11 May 2024 from <https://phys.org/news/2014-09-asian-monsoon-older-previously-thought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.