

## The sound of an atom has been captured

## September 11 2014



On the right, an artificial atom generates sound waves consisting of ripples on the surface of a solid. The sound, known as a surface acoustic wave (SAW) is picked up on the left by a "microphone" composed of interlaced metal fingers. According to theory, the sound consists of a stream of quantum particles, the weakest whisper physically possible. The illustration is not to scale. Credit: Philip Krantz, Krantz NanoArt

Researchers at Chalmers University of Technology are first to show the use of sound to communicate with an artificial atom. They can thereby demonstrate phenomena from quantum physics with sound taking on the role of light. The results will be published in the journal *Science*.

The interaction between atoms and light is well known and has been studied extensively in the field of <u>quantum optics</u>. However, to achieve



the same kind of interaction with <u>sound waves</u> has been a more challenging undertaking. The Chalmers researchers have now succeeded in making acoustic waves couple to an <u>artificial atom</u>. The study was done in collaboration between experimental and theoretical physicists.

"We have opened a new door into the quantum world by talking and listening to atoms", says Per Delsing, head of the experimental research group. "Our long term goal is to harness <u>quantum physics</u> so that we can benefit from its laws, for example in extremely fast computers. We do this by making electrical circuits which obey quantum laws, that we can control and study."

An artificial atom is an example of such a quantum electrical circuit. Just like a regular atom, it can be charged up with energy which it subsequently emits in the form of a particle. This is usually a particle of light, but the atom in the Chalmers experiment is instead designed to both emit and absorb energy in the form of <u>sound</u>.





Microscope image: The artificial atom, in grey-blue to the upper right, can emit and absorb sound that moves across the surface of a microchip. The grey-blue structure to the lower left is the combined loudspeaker/microphone used to communicate acoustically with the atom. Credit: Martin Gustafsson and Maria Ekström

"According to the theory, the sound from the atom is divided into quantum particles", says Martin Gustafsson, the article's first author. "Such a particle is the weakest sound that can be detected."

Since sound moves much slower than light, the acoustic atom opens entire new possibilities for taking control over quantum phenomena.

"Due to the slow speed of sound, we will have time to control the <u>quantum particles</u> while they travel" says Martin Gustafsson. "This is



difficult to achieve with light, which moves 100,000 times more quickly."



Microscope image: This is a zoom-in of the artificial atom, with its integrated Superconducting Quantum Interference Device (SQUID) in violet. The SQUID gives the atom its quantum properties, and the fingers sticking up to the left provide the coupling to sound waves. Credit: Martin Gustafsson and Maria Ekström

The low speed of sound also implies that it has a short wavelength compared to light. An atom that interacts with <u>light</u> waves is always much smaller than the wavelength. However, compared to the wavelength of sound, the atom can be much larger, which means that its properties can be better controlled. For example, one can design the



atom to couple only to certain acoustic frequencies or make the interaction with the sound extremely strong.

The frequency used in the experiment is 4.8 gigahertz, close to the microwave frequencies common in modern wireless networks. In musical terms, this corresponds approximately to a D28, about 20 octaves above the highest note on a grand piano.

At such high frequencies, the wavelength of the sound becomes short enough that it can be guided along the surface of a microchip. On the same chip, the researchers have placed an artificial atom, which is 0.01 millimeters long and made of a superconducting material.

**More information:** "Propagating phonons coupled to an artificial atom" *Science*, 2014. <u>www.sciencemag.org/lookup/doi/ ...</u> <u>1126/science.1257219</u>

The theoretical research group, led by Göran Johansson, recently published a paper on how the acoustic atom functions: <u>link.aps.org/doi/10.1103/PhysRevA.90.013837</u>

Provided by Chalmers University of Technology

Citation: The sound of an atom has been captured (2014, September 11) retrieved 3 May 2024 from <u>https://phys.org/news/2014-09-atom-captured.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.