

Big science from small insects

September 22 2014, by Arran Frood



Researchers working in the RIS lab. Credit: Copyright BBSRC

Anniversaries are often a time to look back. But after taking stock of the past, it can be just as important to look to the future.

2014 saw the fiftieth anniversary of the [Rothamsted Insect Survey](#) (RIS) that has collected and cataloged more than 30M insects across the UK – information that informs farmers of when pest species might hit their crops, what animal disease vectors are flying, and is used by ecologists detecting the patterns that underpin the structure of biological communities.

Yet after 50 years its work is far from done. Indeed, the data become more and more valuable as the time series increases. As a changing climate alters the distribution and abundance of important crop pests, continued monitoring of new insect threats to economically important crops such as cereals is crucial. Many of the insects, principally aphids, also carry viral diseases that further undermine aspects of the UK's food security.

"The Rothamsted Insect Survey has amassed an incredible wealth of data and is now widely regarded as the most comprehensive and continual database in the world on terrestrial invertebrates," says RIS Project Leader [Dr Richard Harrington](#) from Rothamsted Research, an institute that receives strategic funding from BBSRC.

Supporting the survey's work into the 21st century, RIS is now funded as a BBSRC National Capability – an institute-based resource intended to benefit the wider scientific community. It designates the RIS as a repository of data and expertise that can be used by scientists in the UK and across the world.

Indeed, suction-traps of the RIS design are spreading across the world as other countries develop similar surveillance and warning mechanisms, opening opportunities for collaboration between growers and industry with entomologists interested in the fundamental mechanisms behind the spread of insect populations.

Research Impacts of the Rothamsted Insect Survey

Forecasts and pest control

RIS researchers have found strong relationships between winter temperature and the [time that aphids](#) first appear in the suction-traps, and their abundance. These relationships are used to forecast when pest

aphid movement into crops is likely to start, and to predict the impacts of [climate change](#) on pest problems. The wealth of data over time has revealed that compared to 50 years ago, many aphids are flying a month or more earlier. These findings are [disseminated in web](#) and email bulletins that are issued on Fridays, picked up by relevant media, and then used by practitioners to guide aphid control programmes.



A moth light trap. Credit: Copyright Rothamsted Research

Insecticide resistance

Individual aphids are tested for their resistance to a range of insecticides; in particular major pests the peach-potato aphid *Myzus persicae* and English grain aphid *Sitobion avenae*. Molecular tools are used to detect

the genetic mutations conferring resistance, and the findings are fed back to researchers in academia and industry. Recent [analysis at Rothamsted](#) has shown that resistance to pyrethroids is increasing in both England and Scotland.

Plant viruses

Aphids are particularly important [vectors of crop viruses](#), which lower yields. RIS workers detect the presence of certain viruses in aphids and hence their potential as vectors. The traps in Scotland provide data in support of the [Scottish Seed Potato Classification Scheme](#).

Pest insect population dynamics

Analysis of 5715 series of annual moth and aphid species abundance showed, for the first time, the critical importance of RIS' long time series data in the ability to detect [density dependence](#) – an important cornerstone of population dynamic theory that can be used to model and predict insect populations in space and time.

Testing theories of biodiversity



The suction traps are located up and down the UK. Credit: Copyright Google Maps

RIS data were used to test a controversial new theory of the dynamics of biodiversity; this '[Unified neutral theory of biodiversity and biogeography](#)' was based on the assumption that trait differences between trophically similar species (growth rates in aphids for example) had no impact on their relative abundance or speciation rates. Data from RIS' moth light-trap network tested the model but did not fit well because ecological communities fluctuated more than expected.

Aphid winter attrition

The temperature at which aphids die is crucial to their overwintering success – it affects how they will infect crops. Aphids were known to supercool and to freeze at temperatures below -20C, but data from the

suction-trap network suggested much higher temperatures were causing [significant mortality](#) to the mobile forms, probably as a result of membrane disruption.

Effects of climate change

RIS data have been used to detect effects of climate and other environmental changes on the abundance and seasonality of aphids and moths. They have found that, increasingly, [aphids](#) are overwintering in the mobile stages as opposed to laying eggs. Climate change is also implicated in some recent moth declines.

Evolution of plant defences

Europe-wide suction-trap data, combined with laboratory selection experiments, showed that different types of *Arabidopsis thaliana*, a plant much used in lab experiments, are selected according to the relative abundance of [two specialist aphid](#) species. This is significant because it demonstrates that pests can directly affect how their host-plants evolve in relatively short spaces of time.

Provided by Biotechnology and Biological Sciences Research Council

Citation: Big science from small insects (2014, September 22) retrieved 4 May 2024 from <https://phys.org/news/2014-09-big-science-small-insects.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.