


New concepts based on advances in animal systematics

September 17 2014



The new superphylum arrangement is categorized by differences in larval development. Ambulacraria larvae move via cilia while Chordata larvae, through the use of structures like the notochord and backbones are capable of using more powerful muscles to swim through their environment like fish.

The way in which most multicellular organisms have been classified has been the same for more than a century. Only recently have scientists developed the tools and knowledge to question the way we classify organisms. The data accumulated from these newly developed techniques has the potential to change how future generations of scientists classify organisms and understand the connections between

them. Professor Noriyuki Satoh from the Okinawa Institute of Science and Technology Graduate University, as well as Professors Daniel Rokhsar of University California, Berkeley and Teruaki Nishikawa of Toho University, Funabashi, have conducted the research resulting in their new article in Proceedings of the Royal Society B, which proposes a substantial change to how animals, chordates in particular, are classified.

The reasoning behind this proposed reclassification comes from the collection of data from different fields of study, including genomics, evolutionary developmental biology and molecular phylogenetic techniques. Genomics is the broadest of these, where all aspects of the genome of a particular species are analyzed and compared to other genomes to find differences and similarities. Molecular phylogenetic techniques are more specific than genomics, examining both new and old genes from the entire genome and comparing them in groups. Evolutionary developmental biology is the oldest of the methods, which examines an organism's structure or body plan for similarities and differences.

The research proposes that given new data from these methods of observation, there should be a new category called Superphylum. This category would fit between the rank Infrakingdom where Deuterostomia are, and Phylum, which is currently the rank of Chordata. Much of how [organisms](#) are classified depends on their shared characteristics. The number of shared characteristics one organism has with others determines where it is classified.

Deuterostomes include all organisms that follow a very specific embryonic development. Chordates include all animals with a notochord or dorsal neural tube, which are the core components of a central nervous system. Looking at the shared characteristics of organisms from the perspective of evolutionary biology suggests that a larger and more

Provided by Okinawa Institute of Science and Technology

Citation: New concepts based on advances in animal systematics (2014, September 17) retrieved 9 April 2024 from <https://phys.org/news/2014-09-concepts-based-advances-animal-systematics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.