

# Answer to restoring lost island biodiversity found in fossils

September 22 2014

---



A new University of Florida study shows scientists are only beginning to understand the roles of native species in prehistoric island ecosystems. Researchers discovered this 3,000-year-old fossil skull of a Cuban Crocodile, *Crocodylus rhombifer*, in the Bahamas. Credit: Florida Museum of Natural History, by Kristen Grace

Many native species have vanished from tropical islands because of human impact, but University of Florida scientists have discovered how fossils can be used to restore lost biodiversity.

The key lies in organic materials found in fossil bones, which contain evidence for how ancient ecosystems functioned, according to a new study available online and in the September issue of the *Journal of Herpetology*. Pre-human island ecosystems provide vital clues for saving endangered island [species](#) and re-establishing [native species](#), said lead author Alex Hastings, who conducted work for the study as graduate student at the Florida Museum of Natural History and UF department of geological sciences.

"Our work is particularly relevant to [endangered species](#) that are currently living in marginal environments," said Hastings, currently a postdoctoral researcher at Martin Luther University Halle-Wittenberg. "A better understanding of species' natural roles in ecosystems untouched by people might improve their prospects for survival."

Thousands of years ago, the largest carnivore and herbivore on the Bahamian island of Abaco disappeared. The study reconstructs the ancient food web of Abaco where these two mega-reptiles, the endangered Cuban Crocodile (*Crocodylus rhombifer*) and the now-extinct Albury's Tortoise (*Chelonoidis alburyorum*), once flourished. Today, there is no modern terrestrial ecosystem like that of ancient Abaco, with reptiles filling the roles of largest herbivore and carnivore.

In the study, sponsored by the National Science Foundation and National Geographic Society, researchers embarked on the difficult task of reconstructing an ecosystem where few of the components still exist. To understand these missing pieces, scientists analyzed the types of carbon and nitrogen in well-preserved [fossil bones](#) from the Cuban Crocodile and Albury's Tortoise, which was unknown to scientists before its 2004 discovery in the Bahamas. The data reveal the crocodile and tortoise were both terrestrial, showing that reptiles "called the shots" on the island, Hastings said.



Florida Museum of Natural History ornithology curator David Steadman co-authored a new study that reconstructs the ancient food web of the Bahamian island of Abaco where mega-reptiles, including the Cuban Crocodile, once flourished. Credit: Florida Museum of Natural History, by Kristen Grace

The terrestrial nature of these creatures is a great indicator of how biodiversity has changed in the Bahamas and what the ideal circumstances would be for these or similar species to return, said Florida Museum ornithology curator and study co-author David Steadman.

"On islands like Abaco that have always been dominated by reptiles, the flora and fauna are more vulnerable because they have evolved to lead a more laid back, island existence," Steadman said. "Understanding this is important to designing better approaches to conservation on the island."

Early paleontological sites in the Bahamas have yielded bones from numerous species of reptiles, birds and mammals that no longer exist on the islands. James Mead, a vertebrate paleontologist with East Tennessee State University, said more research into the evolutionary history of native plants and animals on Abaco is needed as well as conservation programs based on paleontological research that aims to restore these species.

"The Cuban crocodile is living today in small numbers in Cuba, but this new research shows that it is not living to its fullest potential," Mead said. "The crocodile could live more abundantly in a much wider habitat if we allowed it."

Provided by University of Florida

Citation: Answer to restoring lost island biodiversity found in fossils (2014, September 22) retrieved 20 April 2024 from <https://phys.org/news/2014-09-lost-island-biodiversity-fossils.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.